

HOMOTOPY TYPES OF DIFFEOMORPHISM GROUPS OF NONCOMPACT 2-MANIFOLDS

TATSUHIKO YAGASAKI

ABSTRACT. Suppose M is a noncompact connected smooth 2-manifold without boundary and let $\mathcal{D}(M)_0$ denote the identity component of the diffeomorphism group of M with the compact-open C^∞ -topology. In this paper we investigate the topological type of $\mathcal{D}(M)_0$ and show that $\mathcal{D}(M)_0$ is a topological ℓ_2 -manifold and it has the homotopy type of the circle if M is the plane, the open annulus or the open Möbius band, and it is contractible in all other cases. When M admits a volume form ω , we also discuss the topological type of the group of ω -preserving diffeomorphisms of M . To obtain these results we study some fundamental properties of transformation groups on noncompact spaces endowed with weak topology.

1. INTRODUCTION

The purpose of this paper is the investigation of topological properties of the diffeomorphism groups of noncompact smooth 2-manifolds endowed with the *compact-open* C^∞ -topology. When M is a closed smooth n -manifold, the diffeomorphism group $\mathcal{D}(M)$ with the compact-open C^∞ -topology is a smooth Fréchet manifold [13, Section I.4], and for $n = 2$, S. Smale [26] and C. J. Earle and J. Eell [9] classified the homotopy type of the identity component $\mathcal{D}(M)_0$.

In the C^0 -category, for any compact 2-manifold M , the homeomorphism group $\mathcal{H}(M)$ with the compact-open topology is a topological ℓ_2 -manifold [8, 11, 20, 27], and M. E. Hamstrom [14] classified the homotopy type of the identity component $\mathcal{H}(M)_0$ (cf. [24] for PL-case). In [29] we have shown that $\mathcal{H}(M)_0$ is an ℓ_2 -manifold even if M is a noncompact connected 2-manifold. We also classified its homotopy type and showed that $\mathcal{H}(M)_0$ is contractible except a few cases.

In [2] we studied topological types of transformation groups on noncompact spaces endowed with strong topology. This formulation was intended for an application to homeomorphism groups and diffeomorphism groups of noncompact manifolds endowed with the Whitney topology.

In this article we formulate the notion of weak topology for transformation groups on noncompact spaces. This notion corresponds with the compact-open topology. We see that the main arguments in [28, 29, 30] well extend to transformation groups with weak topology (cf. Theorem 3.1) and these results can be well applied to the diffeomorphism groups of noncompact 2-manifolds.

Suppose M is a smooth n -manifold and X is a closed subset of M . For $r = 1, 2, \dots, \infty$ we denote by $\mathcal{D}_X^r(M)$ the group of C^r -diffeomorphisms h of M onto itself with $h|_X = id_X$, endowed with the compact-open C^r -topology [16, CH.2 Section 1], and by $\mathcal{D}_X^r(M)_0$ the identity connected component of $\mathcal{D}_X^r(M)$. By a compact smooth submanifold of M we mean the union of a disjoint family of a

2000 *Mathematics Subject Classification.* 57S05, 58D05, 57N20, 58A10.

Key words and phrases. Transformation groups, Topological groups, Diffeomorphism groups, Volume-preserving diffeomorphisms, End charge homomorphism, Infinite-dimensional manifolds, σ -compact manifold, Surfaces.

closed smooth k -submanifold of M for $k = 0, 1, \dots, n-1$ and a compact smooth n -submanifold of M . The following is the main result of this paper.

Theorem 1.1. *Suppose M is a noncompact connected smooth 2-manifold without boundary and X is a compact smooth submanifold of M . Then the following hold.*

- (1) $\mathcal{D}_X^r(M)_0$ is a topological ℓ_2 -manifold.
- (2) (i) $\mathcal{D}_X^r(M)_0 \simeq \mathbb{S}^1$ if $(M, X) = (a \text{ plane}, \emptyset), (a \text{ plane}, 1 \text{ pt}), (an \text{ open Möbius band}, \emptyset)$
or $(an \text{ open annulus}, \emptyset)$.
- (ii) $\mathcal{D}_X^r(M)_0 \simeq *$ in all other cases.

Note that any separable infinite-dimensional Fréchet space is homeomorphic to the separable Hilbert space $\ell_2 \equiv \{(x_n) \in \mathbb{R}^\infty : \sum_n x_n^2 < \infty\}$ [6, Chapter VI, Theorem 5.2]. A topological ℓ_2 -manifold is a separable metrizable space which is locally homeomorphic to ℓ_2 . Since topological types of ℓ_2 -manifolds are classified by their homotopy types, Theorem 1.1 (2) implies that $\mathcal{D}_X^r(M)_0 \cong \mathbb{S}^1 \times \ell_2$ in the case (i) and $\mathcal{D}_X^r(M)_0 \cong \ell_2$ in the case (ii).

Let $\mathcal{H}_X(M)_0$ denote the identity connected component of the group of homeomorphisms h of M onto itself with $h|_X = id_X$, endowed with the compact-open C^0 -topology. The comparison of the homotopy types of $\mathcal{D}_X^r(M)_0$ and $\mathcal{H}_X(M)_0$ ([29]) implies the following conclusion:

Corollary 1.1. *Suppose X is a compact smooth submanifold of M . Then the inclusion $\mathcal{D}_X^r(M)_0 \subset \mathcal{H}_X(M)_0$ is a homotopy equivalence.*

For the subgroup of diffeomorphisms with compact supports, we have the following consequences. Let $\mathcal{D}_X^r(M)^c$ denote the subgroup of $\mathcal{D}_X^r(M)$ consisting of diffeomorphisms with compact supports, and let $\mathcal{D}_X^r(M)_0^c$ denote the identity connected component of $\mathcal{D}_X^r(M)^c$. We can also consider the subgroup of diffeomorphisms which are isotopic to id_M by isotopies with compact supports. Let $\mathcal{D}_X^r(M)_0^{c*}$ denote the subgroup of $\mathcal{D}_X^r(M)_0^c$ consisting of $h \in \mathcal{D}_X^r(M)^c$ which admits an ambient C^r -isotopy $h_t : M \rightarrow M \text{ rel } X$ such that $h_0 = h$, $h_1 = id_M$ and h_t ($0 \leq t \leq 1$) have supports in a common compact subset of M .

We say that a subspace A of a space X is homotopy dense (or has the homotopy negligible complement) in X if there exists a homotopy $\varphi_t : X \rightarrow X$ such that $\varphi_0 = id_X$ and $\varphi_t(X) \subset A$ ($0 < t \leq 1$). In this case, the inclusion $A \subset X$ is a (controlled) homotopy equivalence, and when X is metrizable, X is an ANR iff A is an ANR (cf. §2.4).

Theorem 1.2. *Suppose M is a noncompact connected smooth 2-manifold without boundary and X is a compact smooth submanifold of M . Then $\mathcal{D}_X^r(M)_0^{c*}$ is homotopy dense in $\mathcal{D}_X^r(M)_0$*

Corollary 1.2. (1) $\mathcal{D}_X^r(M)_0^c$ and $\mathcal{D}_X^r(M)_0^{c*}$ are ANR's.

- (2) The inclusions $\mathcal{D}_X^r(M)_0^{c*} \subset \mathcal{D}_X^r(M)_0^c \subset \mathcal{D}_X^r(M)_0$ are homotopy equivalences.

We notice that the subgroup $\mathcal{D}_X^r(M)_0^{c*}$ coincides with $\mathcal{D}_X^r(M)_0^c$ except some specific cases.

Proposition 1.1. *Suppose M is a noncompact connected smooth 2-manifold without boundary and X is a compact smooth 2-submanifold of M . Then $\mathcal{D}_X^r(M)_0^{c*} = \mathcal{D}_X^r(M)_0^c$ iff (a) M has no product end or (b) $(M, X) = (a \text{ plane}, \emptyset)$ or $(an \text{ open Möbius band}, \emptyset)$.*

For n -manifolds of finite type we can deduce the following conclusion.

Proposition 1.2. *If $M = \text{Int } N$ for some compact smooth n -manifold N with nonempty boundary and X is a compact smooth submanifold of M , then $\mathcal{D}_X^r(M)_0$ is an ℓ_2 -manifold.*

This proposition is not so obvious, because for the subgroup

$$\mathcal{D}_{\partial N}^r(N)' = \{h \in \mathcal{D}_{\partial N}^r(N) \mid f = \text{id on a neighborhood of } \partial N\}$$

the restriction map $\mathcal{D}_{\partial N}^r(N)' \rightarrow \mathcal{D}^r(\text{Int } N)^c$ is a continuous bijection, but not a homeomorphism.

At this point it is important to compare the above results on the compact-open C^∞ -topology with those on the Whitney C^∞ -topology in [2]. Suppose M is a noncompact connected smooth n -manifold without boundary. Let $\mathcal{D}^\infty(M)_w$ denote the group $\mathcal{D}^\infty(M)$ endowed with the Whitney C^∞ -topology. In [2] we have shown that $\mathcal{D}^\infty(M)_w$ is locally homeomorphic $\square^\omega \ell_2$ (the countable box product of ℓ_2), while $\mathcal{D}^\infty(M)_w^c$ is an $\mathbb{R}^\infty \times \ell_2$ -manifold. Here, $\mathbb{R}^\infty$ is the direct limit of the tower $\mathbb{R}^1 \subset \mathbb{R}^2 \subset \mathbb{R}^3 \subset \dots$. Since the Whitney C^∞ -topology is so strong, it is seen that $(\mathcal{D}^\infty(M)_w)_0 = (\mathcal{D}^\infty(M)_w^c)_0$ and any compact subset of $\mathcal{D}^\infty(M)_w^c$ has a common compact support. When $n = 2$, the difference between $\mathcal{D}^\infty(M)_0$ and $(\mathcal{D}^\infty(M)_w)_0$ is summarized as follows.

Remark 1.1. If M is a noncompact connected smooth 2-manifold without boundary, then

$$(1) \quad \mathcal{D}^\infty(M)_0 \cong \begin{cases} \mathbb{S}^1 \times \ell_2 & \text{if } M = \text{a plane, an open Möbius band, an open annulus,} \\ \ell_2 & \text{in all other cases,} \end{cases}$$

the inclusion $\mathcal{D}^\infty(M)_0^{c*} \subset \mathcal{D}^\infty(M)_0$ is a homotopy equivalence,

$$(2) \quad (\mathcal{D}^\infty(M)_w)_0 = (\mathcal{D}^\infty(M)_w^c)_0 \cong \mathbb{R}^\infty \times \ell_2 \quad ((\mathcal{D}^\infty(M)_w)_0 = \mathcal{D}^\infty(M)_0^{c*} \text{ as sets}).$$

Note that any loop in $(\mathcal{D}^\infty(M)_w)_0$ has a common compact support so that it is inessential, while a loop $(h_t)_t$ in $\mathcal{D}^\infty(M)_0^{c*}$ may not have a common compact support though each h_t admits an isotopy to id_M with compact support. In [30, Example 3.1 (2)] a homotopy equivalence $f : \mathbb{S}^1 \simeq \mathcal{H}(\mathbb{R}^2)_0^{c*}$ is constructed explicitly. This fact can be regarded as a sort of pathology of the compact-open topology.

Next we discuss the groups of volume-preserving diffeomorphisms of noncompact 2-manifolds. Suppose M is a connected oriented smooth n -manifold possibly with boundary and ω is a positive volume form on M . Let $\mathcal{D}^\infty(M; \omega)$ denote the subgroup of $\mathcal{D}^\infty(M)$ consisting of ω -preserving diffeomorphisms of M (endowed with the compact-open C^∞ -topology) and let $\mathcal{D}^\infty(M; \omega)_0$ denote the identity connected component of $\mathcal{D}^\infty(M; \omega)$. In [31, Corollary 1.1] we have shown that the group $\mathcal{D}^\infty(M)_0$ has a factorization

$$(\mathcal{D}^\infty(M)_0, \mathcal{D}^\infty(M; \omega)_0) \cong (\mathcal{V}^+(M; \omega(M), E_M^\omega)_{ew}, \{\omega\}) \times \mathcal{D}^\infty(M; \omega)_0,$$

where $\mathcal{V}^+(M; \omega(M), E_M^\omega)$ is a convex space of positive volume forms on M endowed with the finite-ends weak C^∞ -topology ew (cf. Section 6). This means that the subgroup $\mathcal{D}^\infty(M; \omega)_0$ is a strong deformation retract (SDR) of $\mathcal{D}^\infty(M)_0$. Hence Theorem 1.1 yields the following consequences.

Theorem 1.3. *Suppose M is a noncompact connected orientable smooth 2-manifold without boundary and ω is a volume form on M . Then the following hold.*

$$(1) \quad \mathcal{D}^\infty(M; \omega)_0 \text{ is a topological } \ell_2\text{-manifold and it is a SDR of } \mathcal{D}^\infty(M)_0.$$

- (2) (i) $\mathcal{D}^\infty(M; \omega)_0 \simeq \mathbb{S}^1$ if $M =$ a plane or an open annulus.
(ii) $\mathcal{D}^\infty(M; \omega)_0 \simeq *$ in all other cases.

S. R. Alpern and V. S. Prasad [1] introduced the end charge homomorphism

$$c_0^\omega : \mathcal{D}^\infty(M; \omega)_0 \rightarrow \mathcal{S}(M; \omega).$$

Here $\mathcal{S}(M; \omega)$ is the topological linear space of end charges of (M, ω) and for each $h \in \mathcal{D}^\infty(M; \omega)_0$ the end charge $c_0^\omega(h)$ measures volume transfer toward ends of M under h (cf. Section 6). In [31, Corollary 1.2] we have shown that the homomorphism c^ω has a continuous (non-homomorphic) section. This induces the factorizations

$$(\mathcal{D}^\infty(M; \omega)_0, \ker c_0^\omega) \cong (\mathcal{S}(M; \omega), 0) \times \ker c_0^\omega$$

and implies that the kernel $\ker c_0^\omega$ is a SDR of $\mathcal{D}^\infty(M; \omega)_0$.

Let $\mathcal{D}^\infty(M; \omega)^c$ denote the subgroup of $\mathcal{D}^\infty(M; \omega)$ consisting of ω -preserving diffeomorphisms of M with compact support and let $\mathcal{D}^\infty(M; \omega)_0^c$ denote the identity connected component of $\mathcal{D}^\infty(M; \omega)^c$. The subgroup $\mathcal{D}^\infty(M; \omega)_0^{c*}$ of $\mathcal{D}^\infty(M; \omega)_0^c$ is defined by

$$\mathcal{D}^\infty(M; \omega)_0^{c*} = \{h \in \mathcal{D}^\infty(M; \omega) \mid h \in \mathcal{D}_{M-K}^\infty(M; \omega)_0 \text{ for some compact subset } K \text{ of } M\}.$$

As another application of the results in Section 3, in $n = 2$ we can deduce the topological relations among the subgroups

$$\mathcal{D}^\infty(M; \omega)_0^{c*} \subset \mathcal{D}^\infty(M; \omega)_0^c \subset \ker c_0^\omega.$$

Theorem 1.4. *Suppose M is a noncompact connected orientable smooth 2-manifold without boundary and ω is a volume form on M . Then the following hold.*

- (1) $\ker c_0^\omega$ is an ℓ_2 -manifold and it is a SDR of $\mathcal{D}^\infty(M; \omega)_0$.
- (2) $\mathcal{D}^\infty(M; \omega)_0^{c*}$ is homotopy dense in $\ker c_0^\omega$. Therefore,
 - (i) $\mathcal{D}^\infty(M; \omega)_0^{c*}$ and $\mathcal{D}^\infty(M; \omega)_0^c$ are ANR's, and
 - (ii) the inclusions $\mathcal{D}^\infty(M; \omega)_0^{c*} \subset \mathcal{D}^\infty(M; \omega)_0^c \subset \mathcal{D}^\infty(M; \omega)_0$ are homotopy equivalences.

This paper is organized as follows. Section 2 contains some fundamental facts on ANR's and ℓ_2 -manifolds. Section 3 contains main arguments in this article. Here, we investigate some fundamental topological properties of transformation groups on noncompact spaces endowed with weak topology. In Section 4 these results are applied to the diffeomorphism groups of noncompact manifolds with the compact-open C^r -topology. Theorems 1.1 and 1.2 are proved in Section 5. In Section 6 we discuss the groups of volume-preserving diffeomorphisms of noncompact manifolds.

2. BASIC PROPERTIES OF ANR'S AND ℓ_2 -MANIFOLDS

In Sections 3-6 we see that the ANR-property of diffeomorphism groups and embedding spaces is especially important to investigate the topology of diffeomorphism groups of noncompact manifolds. In this section we recall basic properties of ANR's. We refer to [17, 23] for the theory of ANR's. Throughout the paper we assume that spaces are separable and metrizable and maps are continuous (otherwise specified).

A metrizable space X is called an *ANR* (absolute neighborhood retract) for metrizable spaces if any map $f : B \rightarrow X$ from a closed subset B of a metrizable space Y admits an extension to a neighborhood U of B in Y . If we can always take $U = Y$, then X is called an *AR*. An AR is exactly a contractible ANR. It is well known that X is an AR (an ANR) iff it is a retract of (an open subset of) a normed space. Any ANR has a homotopy type of CW-complex.

We will apply the following criterion of ANR's [17]:

- Lemma 2.1.** (1) *Any retract of an AR (an open subset of an ANR) is an AR (an ANR).*
(2) *A metrizable space X is an ANR iff each point of X has an ANR neighborhood in X .*
(3) *If $X = \cup_{i=1}^{\infty} U_i$, U_i is open in X , $U_i \subset U_{i+1}$ and each U_i is an AR, then X is an AR.*
(4) *$X \times Y$ is a nonempty ANR iff X and Y are nonempty ANR's. In a fiber bundle, the total space is an ANR iff both the base space and the fiber are ANR's.*
(5) *A metric space X is an ANR iff for any $\varepsilon > 0$ there is an ANR Y and maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that gf is ε -homotopic to id_X [15].*

In the statement (3) the space X is an ANR and has the trivial homotopy groups, thus it is contractible. Since any Fréchet space is an AR, by (2) any Fréchet manifold (an ℓ_2 -manifold) is an ANR.

In a fiber bundle, if the base space is contractible and paracompact, then this bundle is trivial. A principal bundle is trivial iff it admits a section. We use the following fact on principal bundles with AR fibers.

Lemma 2.2. *If a principal bundle $p : E \rightarrow B$ has an AR fiber and a metrizable base space B , then,*

- (i) *the bundle p admits a section s and so it is a trivial bundle and*
- (ii) *the map $sp : E \rightarrow E$ is p -fiber-preserving homotopic to id_E and so the map p is a homotopy equivalence with a homotopy inverse s .*

The notion of homotopy denseness (or homotopy negligibility) has been defined in §1. A subspace A of a space X is homotopy dense (HD) in X if there exists a homotopy $\varphi_t : X \rightarrow X$ such that $\varphi_0 = id_X$ and $\varphi_t(X) \subset A$ ($0 < t \leq 1$) ([27]). The homotopy φ_t is called an absorbing homotopy. The map $\varphi_1 : X \rightarrow A$ is a homotopy inverse of the inclusion $A \subset X$.

Lemma 2.3. (1) *Suppose a subspace A is HD in X . Then*

- (i) *the inclusion $A \subset X$ is a (controlled) homotopy equivalence, and*
 - (ii) *X is an ANR iff A is an ANR.*
- (2) *A subspace A is HD in X iff every $x \in X$ admits an open neighborhood U in X and a homotopy $\varphi_t : U \rightarrow X$ such that φ_0 is the inclusion $U \subset X$ and $\varphi_t(U) \subset A$ ($0 < t \leq 1$). (cf. [28, Fact 4.1 (i)]).*

Lemma 2.3(1)(ii) follows from Lemma 2.1(5).

We conclude this preliminary section with the following characterization of ℓ_2 -manifold topological groups [8, 27]. An ℓ_2 -manifold is a separable metrizable space locally homeomorphic to ℓ_2 .

Theorem 2.1. (1) *A topological group is an ℓ_2 -manifold iff it is a separable, non locally compact, completely metrizable ANR.*

(2) *Two ℓ_2 -manifolds are homeomorphic iff they are homotopy equivalent.*

3. TRANSFORMATION GROUPS WITH WEAK TOPOLOGY

This section includes some results on transformation groups with weak topology. In the next section, these results will be applied to the diffeomorphism groups of noncompact manifolds endowed with the compact-open C^r -topology.

3.1. Transformation groups.

A *transformation group* means a pair (G, M) in which M is a locally compact, σ -compact Hausdorff space and G is a topological group acting on M continuously and effectively. Each $g \in G$ induces a homeomorphism of M , which is also denoted by the same symbol g . Let $G^c = \{g \in G \mid \text{supp}(g) \text{ is compact}\}$. For any subsets K, N of M we obtain the following subgroups of G :

$$G_K = \{g \in G : g|_K = \text{id}_K\}, \quad G(N) = G_{M \setminus N}, \quad G_K(N) = G_K \cap G(N), \quad G_K^c = G_K \cap G^c \text{ etc.}$$

For any subgroup H of G , let H_0 denote the connected component of the unit element e in H . For example, the symbol G_0^c denotes the connected component of e in G^c . We also consider a subgroup G_0^{c*} of G_0^c defined by

$$G_0^{c*} = \{h \in G^c \mid h \in G(K)_0 \text{ for some compact subset } K \text{ of } M\}.$$

For subsets $K \subset L \subset N$ of M , we have the set of embeddings

$$\mathcal{E}_K^G(L, N) = \{g|_L : L \rightarrow M \mid g \in G_K(N)\}.$$

(If $K = \emptyset$, the symbol K is omitted from the notation.) The group $G_K(N)$ acts transitively on the set $\mathcal{E}_K^G(L, N)$ by $g \cdot f = gf$ ($g \in G_K(N)$, $f \in \mathcal{E}_K^G(L, N)$). The restriction map

$$r : G_K(N) \rightarrow \mathcal{E}_K^G(L, N), \quad r(g) = g|_L.$$

coincides with the orbit map at the inclusion $i_L : L \subset M$ under this action.

We need to pay an attention on the topology of the space $\mathcal{E}_K^G(L, N)$. A topology on $\mathcal{E}_K^G(L, N)$ is called *admissible* if the $G_K(N)$ -action is continuous with respect to this topology. The strongest admissible topology on $\mathcal{E}_K^G(L, N)$ is the quotient topology induced by the map r . Otherwise specified, the set $\mathcal{E}_K^G(L, N)$ is endowed with this quotient topology. For $K \subset L_1 \subset L \subset N_1 \subset N$, one sees that $\mathcal{E}_K^G(L, N_1)$ is a subspace of $\mathcal{E}_K^G(L, N)$ and the restriction map $\mathcal{E}_K^G(L, N) \rightarrow \mathcal{E}_K^G(L_1, N)$ is continuous.

When the set $\mathcal{E}_K^G(L, N)$ is endowed with a specific admissible topology τ , we write $\mathcal{E}_K^G(L, N)^\tau$ to avoid the ambiguity. The map $r : G_K(N) \rightarrow \mathcal{E}_K^G(L, N)^\tau$ is continuous. For any subset $\mathcal{F}$ of $\mathcal{E}_K^G(L, N)^\tau$, let $\mathcal{F}^\tau$ denote the space $\mathcal{F}$ endowed with the subspace topology induced from τ . When $i_L \in \mathcal{F}$, let $\mathcal{F}_0^\tau$ denote the connected component of i_L in $\mathcal{F}^\tau$.

We say that a map $f : X \rightarrow Y$ has a local section at $y \in Y$ if there exists a neighborhood U of y in Y and a map $s : U \rightarrow X$ with $fs = i_U$.

Lemma 3.1. *Suppose H is a subgroup of $G_K(N)$ and the restriction map $r : G_K(N) \rightarrow \mathcal{E}_K^G(L, N)^\tau$ has a local section $s : \mathcal{U} \rightarrow H \subset G_K(N)$ at i_L . Then, the following hold.*

- (1) *The restriction map $r|_H : H \rightarrow \mathcal{E}^H(L, N)^\tau$ is a principal bundle with the structure group H_L .*
- (2) (i) *The topology τ coincides with the quotient topology.*
(ii) *$\mathcal{E}^H(L, N)^\tau$ is open in $\mathcal{E}_K^G(L, N)^\tau$.*
(iii) *If H is a normal subgroup of $G_K(N)$, then*
 - (a) *each orbit of H is closed and open in $\mathcal{E}_K^G(L, N)^\tau$*
(in particular, $\mathcal{E}^H(L, N)^\tau$ is closed and open in $\mathcal{E}_K^G(L, N)^\tau$),
 - (b) *if $H \subset G_K(N)_0$, then $\mathcal{E}^H(L, N)^\tau = \mathcal{E}_K^G(L, N)_0^\tau$.*

Proof. (1) Note that (a) $\mathcal{E}^H(L, N)^\tau = H \cdot i_L$ and the map $r|_H$ coincides with the orbit map at i_L under the action of the group H and (b) $\mathcal{U} \subset \mathcal{E}^H(L, N)^\tau$ and s is a local section of $r|_H$ at i_L . Hence, the statement (1) follows from the well known fact on the orbit map in the theory of group action.

(2) (i) By (1) the map r itself is a principal bundle. Hence, the map r is a quotient map and the topology τ coincides with the quotient topology.

(ii) We may assume that $\mathcal{U}$ is open in $\mathcal{E}_K^G(L, N)^\tau$. Since $\mathcal{U} \subset \mathcal{E}^H(L, N)^\tau$, we have $\mathcal{E}^H(L, N)^\tau = H\mathcal{U}$, which is open in $\mathcal{E}_K^G(L, N)^\tau$.

(iii) Since Hi_L is open, the orbit $H(g \cdot i_L) = g(Hi_L)$ is also open in $\mathcal{E}_K^G(L, N)^\tau$. □

In many cases, the set $\mathcal{E}_K^G(L, N)$ admits a natural admissible topology τ (for instance, the compact-open C^r -topology ($r = 0, 1, \dots, \infty$)). However, this topology τ coincides with the quotient topology, once we obtain a bundle theorem under the topology τ . Hence, our convention does not lose a generality of the arguments in the subsections below.

Finally we extract a behavior of the compact-open C^∞ topology on diffeomorphism groups of a noncompact manifold M at the ends of M extend the notion of the compact-open C^∞ topology on diffeomorphism groups to transformation groups.

Definition 3.1. We say that a transformation group (G, M) has a *weak* topology if it satisfies the following condition:

- (*) For any neighborhood U of e in G there exists a compact subset K of M such that $G_K \subset U$.

Remark 3.1. (1) If G admits a compatible metric ρ , then (G, M) has a weak topology iff for any $\varepsilon > 0$ there exists a compact subset K of M such that $\text{diam}_\rho G_K < \varepsilon$. We can always take ρ to be left-invariant.

(2) If a transformation group (G, M) has a weak topology, then so is (H, M) for any subgroup H of G .

The notion of weak topology is an extension of the notion of compact-open topology to transformation groups. It is readily seen that the compact-open C^∞ topology on the diffeomorphism groups of smooth manifolds is also an example of weak topology.

3.2. Basic assumptions on transformation groups.

Suppose (G, M) is a transformation group. Since M is locally compact and σ -compact, there exists a sequence $\{M_i\}_{i \geq 1}$ of compact subsets of M such that

$$M = \bigcup_{i=0}^{\infty} M_i \quad \text{and} \quad M_{i-1} \subset \text{Int}_M M_i \quad (i \geq 1), \quad \text{where } M_0 = \emptyset.$$

This sequence is called an *exhausting sequence* of M . Let $U_i = \text{Int}_M M_i$ ($i \geq 1$). Consider the following basic conditions on the tuple $(G, M, \{M_i\}_{i \geq 1})$.

Assumption (A).

(A-0) The group G is metrizable and the transformation group (G, M) has a weak topology.

(A-1) For each $j > i > k \geq 0$, the restriction map

$$\pi_{k,j}^i : G_{M_k}(U_j)_0 \longrightarrow \mathcal{E}_{M_k}^G(M_i, U_j)_0$$

is a principal bundle with the structure group $\mathcal{G}_{k,j}^i \equiv G_{M_k}(U_j)_0 \cap G_{M_i}$.

(A-2) (i) $G(U_i)_0$ is an ANR for each $i \geq 1$.

(ii) The spaces $\mathcal{U}_{k,j}^i = \mathcal{E}_{M_k}^G(M_i, U_j)_0$ ($j > i > k \geq 0$) satisfy the next conditions:

$\mathcal{U}_{k,j}^i$ is an open subspace of $\mathcal{E}_{M_k}^G(M_i, M)_0$, $\mathcal{E}_{M_k}^G(M_i, M)_0 = \bigcup_{j>i} \mathcal{U}_{k,j}^i$ and $cl \mathcal{U}_{k,j}^i \subset \mathcal{U}_{k,j+1}^i$.

Below we assume that the tuple $(M, G, \{M_i\}_{i \geq 1})$ satisfies the assumption (A).

Lemma 3.2. (1) For each $i > k \geq 0$, the restriction map

$$\pi_k^i : (G_{M_k})_0 \longrightarrow \mathcal{E}_{M_k}^G(M_i, M)_0$$

is a principal bundle with the structure group $\mathcal{G}_k^i \equiv (G_{M_k})_0 \cap G_{M_i}$.

(2) The spaces $G_{M_k}(U_i)_0$ and $\mathcal{E}_{M_k}^G(M_i, M)_0$ are ANR's for each $i > k \geq 0$.

Proof. (1) This lemma follows from (A-1), (A-2)(ii) and Lemma 3.1.

(2) (i) By (A-1), for each $i > k \geq 1$, the restriction map

$$\pi_{0,i}^k : G(U_i)_0 \longrightarrow \mathcal{E}^G(M_k, U_i)_0$$

is a principal bundle with the fiber $\mathcal{G}_{0,i}^k \equiv G(U_i)_0 \cap G_{M_k}$. Since $G(U_i)_0$ is an ANR by (A-2)(i), so are $\mathcal{G}_{0,i}^k$ and $G_{M_k}(U_i)_0 = (\mathcal{G}_{0,i}^k)_0$.

(ii) By (i) and (A-1) $\mathcal{E}_{M_k}^G(M_i, U_j)_0$ is an ANR for each $j > i$, and so is $\mathcal{E}_{M_k}^G(M_i, M)_0$ by (A-2)(ii). $\square$

Lemma 3.3. (1)(i) If G_0 is an ANR, then so is $(G_{M_i})_0$ for any $i \geq 1$.

(ii) If G_{M_i} is an ANR for some $i \geq 1$, then so is G_0 .

(2) If $(G_{M_i})_0^{c*}$ is HD in $(G_{M_i})_0$ and $(G_{M_i})_0$ is open in G_{M_i} for some $i \geq 1$, then G_0^{c*} is HD in G_0 .

Proof. (1) The restriction map $\pi_0^i : G_0 \rightarrow \mathcal{E}^G(M_i, M)_0$ is a bundle map with fiber $\mathcal{G}_0^i = G_0 \cap G_{M_i}$. Hence, there exists an open neighborhood $\mathcal{U}$ of the inclusion map i_{M_i} in $\mathcal{E}^G(M_i, M)_0$ such that $(\pi_0^i)^{-1}(\mathcal{U}) \cong \mathcal{U} \times \mathcal{G}_0^i$. Since $\mathcal{U}$ is an ANR by (A-1)(i), it follows that $(\pi_0^i)^{-1}(\mathcal{U})$ is an ANR if and only if $\mathcal{G}_0^i$ is an ANR. The assertions (i) and (ii) follow from these observations.

(2) By Lemma 2.3 (2) it suffices to verify the following assertion:

(#) Every $h \in G_0$ admits an open neighborhood $\mathcal{V}$ in G_0 and a homotopy $\varphi : \mathcal{V} \times [0, 1] \rightarrow G_0$ such that φ_0 is the inclusion $\mathcal{V} \subset G_0$ and $\varphi_t(\mathcal{V}) \subset G_0^{c*}$ ($0 < t \leq 1$).

Under the map $\pi_0^i : G_0 \rightarrow \mathcal{E}^G(M_i, M)_0$, each $h \in G_0$ induces $h|_{M_i} \in \mathcal{E}^G(M_i, M)_0$, and by (A-1)(ii) we have $h|_{M_i} \in \mathcal{E}^G(M_i, U_j)_0$ for some $j > i$. By (A-2) the restriction map

$$\pi_{0,j}^i : G(U_j)_0 \longrightarrow \mathcal{E}^G(M_i, U_j)_0$$

is a bundle map and so there exists an open neighborhood $\mathcal{U}$ of $h|_N$ in $\mathcal{E}^G(M_i, U_j)_0$ and a local section $s : \mathcal{U} \rightarrow G(U_j)_0$ of $\pi_{0,j}^i$ such that $s(h|_{M_i}) = h$. Choose a small open neighborhood $\mathcal{V}$ of h in G_0 such that $\pi_0^i(\mathcal{V}) \subset \mathcal{U}$. For each $g \in \mathcal{V}$ we have $s(g|_{M_i})|_{M_i} = g|_{M_i}$ and $s(g|_{M_i})^{-1}g \in G_{M_i}$. Since $s(h|_{M_i})^{-1}h = id_M \in (G_{M_i})_0$ and $(G_{M_i})_0$ is open in G_{M_i} , by replacing $\mathcal{V}$ by a smaller one, we may assume that $s(g|_{M_i})^{-1}g \in (G_{M_i})_0$ ($g \in \mathcal{V}$).

There exists an absorbing homotopy ψ_t of $(G_{M_i})_0$ into $(G_{M_i})_0^{c*}$. Then the required homotopy $\varphi_t : \mathcal{V} \rightarrow G_0$ is defined by

$$\varphi_t(g) = s(g|_{M_i})\psi_t(s(g|_{M_i})^{-1}g).$$

Then $\varphi_0(g) = g$ and for $0 < t \leq 1$ we have $\varphi_t(g) \in G_0^{c*}$ since $s(g|_{M_i}) \in G(U_j)_0 \subset G_0^{c*}$ and $\psi_t(s(g|_{M_i})^{-1}g) \in (G_{M_i})_0^{c*} \subset G_0^{c*}$. $\square$

3.3. Contractibility conditions.

In this subsection we deduce some conclusions under some contractibility conditions. Consider the following conditions on the tuple $(M, G, \{M_i\}_{i \geq 1})$:

Condition (C).

(C-1) $\mathcal{E}_{M_k}^G(M_i, M)_0 \simeq *$ for each $i > k \geq 0$.

(C-2) $\mathcal{G}_{0,j}^i \simeq *$ for each $j > i \geq 1$.

Below we assume that the tuple $(G, M, \{M_i\}_{i \geq 1})$ satisfies the assumption (A). Since G is metrizable by the condition (A-0), it admits a left-invariant metric ρ .

Lemma 3.4. *If $(G, M, \{M_i\}_{i \geq 1})$ satisfies the condition (C-1), then the following hold.*

- (1) For each $i > k \geq 0$,
 - (i) the bundle π_k^i is a trivial bundle and $\mathcal{G}_k^i = (G_{M_i})_0$,
 - (ii) $(G_{M_k})_0$ strongly deformation retracts onto $(G_{M_i})_0$.
- (2) $(G_{M_k})_0$ is an AR for each $k \geq 0$.

Proof. The conditions (A-1) and (C-1) imply that $\mathcal{E}_{M_k}^G(M_i, M)_0$ is an AR.

(1) Since the base space $\mathcal{E}_{M_k}^G(M_i, M)_0$ is metrizable and contractible, the bundle π_k^i is trivial. This means that there exists a fiber-preserving homeomorphism over $\mathcal{E}_{M_k}^G(M_i, M)_0$,

$$(G_{M_k})_0 \cong \mathcal{E}_{M_k}^G(M_i, M)_0 \times \mathcal{G}_k^i.$$

Since $(G_{M_k})_0$ is connected, so is $\mathcal{G}_k^i$ and we have $\mathcal{G}_k^i = (G_{M_i})_0$. Since the base $\mathcal{E}_{M_k}^G(M_i, M)_0$ is an AR, it admits a strong deformation retraction (SDR) onto the singleton $\{i_{M_i}\}$. This induces the required SDR of $(G_{M_k})_0$ onto $\mathcal{G}_k^i$

(2)(i) First we show that $(G_{M_k})_0 \simeq *$. By (1)(ii), for each $i \geq 0$ there exists a SDR h_t^i ($0 \leq t \leq 1$) of $(G_{M_i})_0$ onto $(G_{M_{i+1}})_0$. A SDR h_t ($k \leq t \leq \infty$) of $(G_{M_k})_0$ onto $\{id_M\}$ is defined by

$$\begin{aligned} h_t(f) &= h_{t-i}^i h_1^{i-1} \cdots h_1^k(f) \quad (f \in (G_{M_k})_0, i \geq k, i \leq t \leq i+1) \\ h_\infty(f) &= id_M. \end{aligned}$$

Since (G, M) has a weak topology, it follows that $\text{diam}_\rho(G_{M_i})_0 \rightarrow 0$ and the homotopy $h : (G_{M_k})_0 \times [k, \infty] \rightarrow (G_{M_k})_0$ is continuous.

(ii) To see that $(G_{M_k})_0$ is an ANR, we apply Lemma 2.1 (5) (the Hanner's criterion). By (1), for each $i > k$ the restriction map

$$\pi_k^i : (G_{M_k})_0 \longrightarrow \mathcal{E}_{M_k}^G(M_i, M)_0$$

is a trivial bundle with an ANR base space and the fiber $\mathcal{G}_k^i \equiv (G_{M_i})_0$. By (i) the fiber $\mathcal{G}_k^i$ is contractible. These imply that π_k^i admits a section s_k^i and that $s_k^i \pi_k^i$ is π_k^i -fiber preserving homotopic to id . Since the metric ρ is left-invariant, the diameter of each fiber of π_k^i coincides with $\text{diam}_\rho G_{M_i}$. Since $\text{diam}_\rho G_{M_i} \rightarrow 0$ ($i \rightarrow 0$), the Hanner's criterion implies that $(G_{M_k})_0$ is an ANR. $\square$

Lemma 3.5. *If $(G, M, \{M_i\}_{i \geq 1})$ satisfies the condition (C-2), then G_0^{c*} is HD in G_0 .*

Proof. (1) For notational simplicity, for each $j > i \geq 1$ we set $\mathcal{D}_j = G(U_j)_0$ and $\mathcal{U}_{i,j} = \mathcal{E}^G(M_i, U_j)_0$. By (A-2) the map $\pi_{0,j}^i : \mathcal{D}_j \rightarrow \mathcal{U}_{i,j}$ is a principal bundle. Since the fiber is an AR by (C-2) and the base space is metrizable, this bundle has a global section. Thus, the map $\pi_{0,j}^i$ is a trivial bundle with an AR fiber and hence it has the following relative lifting property:

(*) If Y is a metric space, B is a closed subset of Y and $\varphi : Y \rightarrow \mathcal{U}_{i,j}$ and $\varphi_0 : B \rightarrow \mathcal{D}_j$ are maps with $\varphi|_B = \pi_{0,j}^i \varphi_0$, then there exists a map $\Phi : Y \rightarrow \mathcal{D}_j$ such that $\pi_{0,j}^i \Phi = \varphi$ and $\Phi|_B = \varphi_0$.

(2) Next consider the principal bundle

$$\pi_0^i : G_0 \rightarrow \mathcal{E}^G(M_i, M)_0.$$

For each $j > i \geq 1$, we set $\mathcal{V}_{i,j} = (\pi_0^i)^{-1}(\mathcal{U}_{i,j}) \subset G_0$. Then $\mathcal{U}_{i,j}$, $\mathcal{V}_{i,j}$ and $\mathcal{D}_j$ satisfy the following conditions: for each $i \geq 1$

- (i) $\mathcal{E}^G(M_i, M)_0 = \cup_{j>i} \mathcal{U}_{i,j}$, $\mathcal{U}_{i,j}$ is open in $\mathcal{E}^G(M_i, M)_0$ and $cl \mathcal{U}_{i,j} \subset \mathcal{U}_{i,j+1}$.
- (ii) $G_0 = \cup_{j>i} \mathcal{V}_{i,j}$, $\mathcal{V}_{i,j}$ is open in G_0 , $cl \mathcal{V}_{i,j} \subset \mathcal{V}_{i,j+1}$ and $\mathcal{V}_{i+1,j} \subset \mathcal{V}_{i,j}$ ($j > i+1$).
- (iii) $G_0^{c*} = \cup_{j>i} \mathcal{D}_j$ and $\mathcal{D}_j \subset \mathcal{D}_{j+1}$.

(3) We have to construct a homotopy $F : G_0 \times [0, 1] \rightarrow G_0$ such that $F_0 = id$ and $F_t(G_0) \subset G_0^{c*}$ ($0 < t \leq 1$). We replace the interval $[0, 1]$ by $[1, \infty]$.

(i) For each $i \geq 1$ we can find a map $s^i : \mathcal{E}^G(M_i, M)_0 \rightarrow G_0^{c*}$ such that

$$s^i(f)|_{M_i} = f|_{M_i} \quad (f \in \mathcal{E}^G(M_i, M)_0) \text{ and } s^i(cl \mathcal{U}_{i,j}) \subset \mathcal{D}_{j+1} \quad (j > i).$$

In fact, using the property (*), inductively we can construct maps $s_j^i : cl \mathcal{U}_{i,j} \rightarrow \mathcal{D}_{j+1}$ ($j > i$) such that

$$s_j^i(f)|_{M_i} = f|_{M_i} \quad (f \in cl \mathcal{U}_{i,j}) \text{ and } s_{j+1}^i|_{cl \mathcal{U}_{i,j}} = s_j^i.$$

The map s^i is defined by $s^i|_{cl \mathcal{U}_{i,j}} = s_j^i$. Let $F_i = s^i \pi_0^i : G_0 \rightarrow G_0^{c*}$. We have $F_i(cl \mathcal{V}_{i,j}) \subset \mathcal{D}_{j+1}$ and $F_i(h)|_{M_i} = h|_{M_i}$.

(ii) For each $i \geq 1$, we can inductively construct a sequence of homotopies

$$H^j : cl \mathcal{V}_{i+1,j} \times [i, i+1] \rightarrow \mathcal{D}_{j+1} \quad (j > i+1)$$

such that

$$H_i^j = F_i, \quad H_{i+1}^j = F_{i+1}, \quad H^{j+1}|_{cl \mathcal{V}_{i+1,j} \times [i, i+1]} = H^j \quad \text{and} \quad H_t^j(h)|_{M_i} = h|_{M_i}.$$

If H^j is given, then H^{j+1} is obtained by applying the property (*) to the diagram:

$$\begin{array}{ccc} B & \xrightarrow{\varphi_0} & \mathcal{D}_{j+2} \\ \cap & & \downarrow \\ Y & \xrightarrow{\varphi} & \mathcal{U}_{i,j+2}, \end{array} \quad \varphi(h, t) = h|_{M_i}, \quad \varphi_0(h, t) = \begin{cases} H^j(h, t) & (h \in cl \mathcal{V}_{i+1,j}) \\ F_t(h) & (t = i, i+1) \end{cases}$$

$$(Y, B) = (cl \mathcal{V}_{i+1,j+1} \times [i, i+1], (cl \mathcal{V}_{i+1,j} \times [i, i+1]) \cup (cl \mathcal{V}_{i+1,j+1} \times \{i, i+1\})).$$

Thus we can define a homotopy $F : G_0 \times [i, i+1] \rightarrow G_0^{c*}$ by $F = H^j$ on $cl \mathcal{V}_{i+1,j} \times [i, i+1]$. Since $F_t(h)|_{M_i} = h|_{M_i}$ for $t \geq i$, we can continuously extend F by $F_\infty = id$. $\square$

Lemmas 3.4 and 3.5 yield the following criterions.

Theorem 3.1. *Suppose the tuple $(G, M, \{M_i\}_{i \geq 1})$ satisfies the assumption (A).*

- (1) *If $G_{M_k}(U_i)_0 \simeq *$ and $\mathcal{G}_{k,j}^i$ is connected (i.e., $G_{M_k}(U_j)_0 \cap G_{M_i} = G_{M_i}(U_j)_0$) for each $j > i > k \geq 0$, then G_0 is an AR and G_0^{c*} is HD in G_0 .*
- (2) (i) *If $G_{M_i} \simeq *$ for each $i \geq 1$, then G_0 is an ANR.*
(ii) *If G_{M_1} is connected and $\mathcal{G}_{1,j}^i = G_{M_1}(U_j)_0 \cap G_{M_i} \simeq *$ for each $j > i \geq 2$, then G_0^{c*} is HD in G_0 .*

Proof. (1) By Lemmas 3.4 and 3.5 it suffices to verify the following assertions: for each $j > i > k \geq 0$

- (i) the bundle $\pi_{k,j}^i$ is trivial, and the spaces $\mathcal{G}_{k,j}^i$, $G_{M_k}(U_j)_0$ and $\mathcal{E}_{M_k}^G(M_i, U_j)_0$ are AR's,
- (ii) $\mathcal{E}_{M_k}^G(M_i, M)_0$ is an AR.

(i) By Lemma 3.2 (2) and the assumption, both the total space $G_{M_k}(U_j)_0$ and the fiber $\mathcal{G}_{k,j}^i = G_{M_i}(U_j)_0$ are AR's. From the homotopy exact sequence of the bundle $\pi_{k,j}^i$ it follows that the ANR base space $\mathcal{E}_{M_k}^G(M_i, U_j)_0$ is contractible and the bundle $\pi_{k,j}^i$ is trivial. (Alternatively, since the fiber is an AR, the principal bundle $\pi_{k,j}^i$ admits a global section, which means that this bundle is trivial and the base space is an AR since it is a retract of the AR total space.)

(ii) The assertion follows from (i), (A-2)(ii) and Lemma 2.1 (3).

(2) Since $(G_{M_1})_0 = G_{M_1}$, by Lemma 3.3 it suffices to show the next assertions:

- (i') $(G_{M_1})_0$ is an ANR if $G_{M_i} \simeq *$ for each $i \geq 1$.
- (ii') $(G_{M_1})_0^{c*}$ is HD in $(G_{M_1})_0$ if $\mathcal{G}_{1,j}^i = G_{M_1}(U_j)_0 \cap G_{M_i} \simeq *$ for each $j > i \geq 2$.

Since the tuple $(G_{M_1}, M, \{M_i\}_{i \geq 2})$ also satisfies the assumption (A), by Lemmas 3.4 and 3.5 it remains to verify the condition (C-1) in (i') and (C-2) in (ii') respectively.

(i') $\mathcal{E}_{M_k}^G(M_i, M)_0 \simeq *$ for each $i > k \geq 1$: In fact, the restriction map

$$\pi_k^i : (G_{M_k})_0 \longrightarrow \mathcal{E}_{M_k}^G(M_i, M)_0$$

is a principal bundle with the fiber $\mathcal{G}_k^i \equiv (G_{M_k})_0 \cap G_{M_i}$. Since $(G_{M_k})_0 = G_{M_k} \simeq *$ and $\mathcal{G}_k^i = G_{M_i} \simeq *$, it follows that the ANR base space $\mathcal{E}_{M_k}^G(M_i, M)_0$ is also contractible.

(ii') $\mathcal{G}_{1,j}^i = G_{M_1}(U_j)_0 \cap G_{M_i} \simeq *$ for each $j > i \geq 2$: This is the condition (C-2) itself for the tuple $(G_{M_1}, M, \{M_i\}_{i \geq 2})$. This completes the proof. $\square$

4. DIFFEOMORPHISM GROUPS OF NON-COMPACT n -MANIFOLDS

4.1. General properties of diffeomorphism groups of n -manifolds.

Suppose M is a smooth (separable metrizable) n -manifold without boundary and X is a closed subset of M with $X \neq M$. Let $r = 1, 2, \dots, \infty$.

When N is a smooth manifold, the symbol $\mathcal{C}^r(M, N)$ denotes the space of C^r -maps $f : M \rightarrow N$ with the compact-open C^r -topology. For any map $f_0 : X \rightarrow N$, let $\mathcal{C}_{f_0}^r(M, N)$ denote the subspace $\{f \in \mathcal{C}^r(M, N) \mid f|_X = f_0\}$. When M is a smooth submanifold of N , the symbol $\mathcal{E}_X^r(M, N)$ denotes the subspace of $\mathcal{C}^r(M, N)$ consisting of C^r -embeddings $f : M \hookrightarrow N$ with $f|_X = id_X$ and $\mathcal{E}_X^r(M, N)_0$ denotes the connected component of the inclusion $i_N : N \subset M$ in $\mathcal{E}_X^r(N, M)$. For spaces Y and Z , the symbol $\mathcal{C}^0(Y, Z)$ denotes the space of C^0 -maps $f : Y \rightarrow Z$ with the compact-open topology.

Consider the jet-map $j^r : \mathcal{C}^r(M, N) \rightarrow \mathcal{C}^0(M, J^r(M, N))$. It is a closed embedding [16, Ch2. Section 4, p.61–62]. Thus, if we choose a complete metric d on the jet-bundle $J^r(M, N)$ with $d \leq 1$ and a sequence of compact n -submanifolds M_i ($i \geq 1$) such that $M_i \subset \text{Int } M_{i+1}$ and $M = \cup_i M_i$, then we can define a complete metric ρ on $\mathcal{C}^r(M, N)$ by

$$\rho(f, g) = \sum_{i=1}^{\infty} \frac{1}{2^i} \sup_{x \in M_i} d(j_x^r f, j_x^r g).$$

When $N = M$, it induces a metric on $\mathcal{D}_X^r(M) \subset \mathcal{C}^r(M, M)$. We can define a complete metric ρ^* on $\mathcal{D}_X^r(M)$ by

$$\rho^*(f, g) = \rho(f, g) + \rho(f^{-1}, g^{-1}).$$

Lemma 4.1. (1) $\mathcal{D}_X^r(M)$ is a topological group [16, Ch 2., Section 4.p. 64].

(2) $\mathcal{D}_X^r(M)$ and $\mathcal{D}_X^r(M)_0$ are separable, completely metrizable and not locally compact [16, Ch 2., Section 4, p. 61–62, Theorems 4.3, 4.4].

(3) $\mathcal{D}_X^r(M)$ and $\mathcal{D}_X^r(M)_0$ are ℓ^2 -manifolds iff they are ANR's.

If M is a closed smooth n -manifold, then $\mathcal{D}^r(M)$ is a smooth Fréchet manifold [13, Section I.4, Example 4.1.3, etc.]. In this paper we are only concerned with topological Fréchet manifolds (= topological ℓ_2 -manifolds). Below, the emphasis is put on relative cases.

Suppose M is a compact smooth n -manifold (possibly with boundary) and X is a closed subset of M . When $E \rightarrow M$ is a smooth fiber bundle over M and $s_0 : X \rightarrow E$ is a section of E over X , the symbol $\Gamma_{s_0}^r(M, E)$ denotes the space of C^r -sections s of E over M with $s|_X = s_0$, endowed with the compact-open C^r -topology. When E is a vector bundle over M , the space $\Gamma_{s_0}^r(M, E)$ is a Fréchet space. If E is a fiber bundle over M and the fiber of E is a smooth manifold without boundary, then $\Gamma_{s_0}^r(M, E)$ is a Fréchet manifold. A local chart around $s \in \Gamma_{s_0}^r(M, E)$ is defined as follows: Consider the restriction $T^v(E)|_{s(M)}$ of the vertical tangent bundle $T^v(E)$ of E to the image $s(M)$ in E . There

exists a diffeomorphism between an open neighborhood of $s(M)$ in E and an open neighborhood of the image of zero section 0 in $T^v(E)|_{s(M)}$. It induces a homeomorphism between an open neighborhood of s in $\Gamma_{s_0}^r(M, E)$ and an open neighborhood of 0 in $\Gamma_{0_X}^r(s(M), T^v(E)|_{s(M)})$. If N is a smooth manifold without boundary and $f_0 : X \rightarrow N$ is a map, then the space $\mathcal{C}_{f_0}^r(M, N) = \{f \in \mathcal{C}^r(M, N) \mid f|_X = f_0\}$ is also a Fréchet manifold since it is identified with $\Gamma_{s_0}^r(M, E)$ for the trivial bundle $E = M \times N \rightarrow M$ and the section $s_0(x) = (x, f_0(x))$ over X .

Lemma 4.2. (i) Suppose N is a smooth manifold without boundary, M is a compact smooth submanifold of N and X is a closed subset of M . Then $\mathcal{E}_X^r(M, N)$ is a Fréchet manifold.

(ii) Suppose M is a compact smooth n -manifold and X is a closed subset of M . If $\partial M \subset X$ or $\partial M \cap X = \emptyset$, then $\mathcal{D}_X^r(M)$ is a Fréchet manifold.

Proof. (i) $\mathcal{E}_X^r(M, N)$ is open in $\mathcal{C}_{i_X}^r(M, N)$ [16, Ch.2 Theorem 1.4].

(ii) If $\partial M \subset X$, then $\mathcal{D}_X^r(M) = \mathcal{E}_X^r(M, M) = \mathcal{E}_X^r(M, \tilde{M})$, where $\tilde{M}$ is the open manifold obtained from M by attaching an open collar to ∂M . In the case $X \cap \partial M = \emptyset$, consider the restriction map $\pi : \mathcal{D}_X^r(M) \rightarrow \mathcal{D}^r(\partial M)$, $\pi(h) = h|_{\partial M}$. Using a collar of ∂M in M , for any $f \in \mathcal{D}^r(\partial M)$ which is sufficiently close to id , we can easily construct a canonical extension $\tilde{f} \in \mathcal{D}_X^r(M)$ of f . This implies that $\text{Im } \pi$ is a clopen subset of $\mathcal{D}^r(\partial M)$ and $\mathcal{D}_X^r(M) \rightarrow \text{Im } \pi$ is a principal bundle with fiber $\mathcal{D}_{X \cup \partial M}^r(M)$. Since $\mathcal{D}^r(\partial M)$ and $\mathcal{D}_{X \cup \partial M}^r(M)$ are Fréchet manifolds, so is $\mathcal{D}_X^r(M)$. $\square$

Remark 4.1. (1) A family $h_t \in \mathcal{D}_X^r(M)$ ($t \in [0, 1]$) is called a C^r -isotopy rel X if $H : M \times [0, 1] \rightarrow M$, $H(x, t) = h_t(x)$, is a C^r -map. In this case h_t is a path in $\mathcal{D}_X^r(M)$ (i.e., $[0, 1] \ni t \mapsto h_t \in \mathcal{D}_X^r(M)$ is continuous). In general, if h_t is a path in $\mathcal{D}_X^r(M)$, then h_t is a C^0 -isotopy rel X , but H is not necessarily C^1 in t .

(2) Since Fréchet manifolds are locally path-connected, the connected components $\mathcal{E}_X^r(N, M)_0$ and $\mathcal{D}_X^r(M)_0$ in Lemma 2.2 are path-connected. Thus, if $h \in \mathcal{D}_X^r(M)_0$, then there is a path h_t ($t \in [0, 1]$) in $\mathcal{D}_X^r(M)_0$ with $h_0 = id_M$ and $h_1 = h$, which is a C^0 -isotopy rel X .

4.2. The bundle theorem.

The bundle theorem connecting diffeomorphism groups and embedding spaces [7, 18, 22, 25] plays an essential role in order to apply Theorem 3.1 (recall Assumption (A)). Suppose M is a smooth n -manifold without boundary. A compact smooth submanifold of M means a compact subset N of M which is the union of a disjoint family $\{N_k\}_{k=0}^n$ such that N_k is a (possibly empty) closed smooth k -submanifold of M for $k = 0, 1, \dots, n-1$ and N_n is a (possibly empty) compact smooth n -submanifold of M .

Suppose N is a compact smooth submanifold of M and X is a closed subset of N . Let U be any open neighborhood of N in M .

Theorem 4.1. For any $f \in \mathcal{E}_X^r(N, U)$ there exist a neighborhood $\mathcal{U}$ of f in $\mathcal{E}_X^r(N, U)$ and a map $\varphi : \mathcal{U} \rightarrow \mathcal{D}_{X \cup (M \setminus U)}^r(M)_0$ such that $\varphi(g)f = g$ ($g \in \mathcal{U}$) and $\varphi(f) = id_M$.

Consider the restriction map $\pi : \mathcal{D}_{X \cup (M \setminus U)}^r(M) \rightarrow \mathcal{E}_X^r(N, U)$, $\pi(h) = h|_N$. The group $\mathcal{D}_{N \cup (M \setminus U)}^r(M)$ acts on $\mathcal{D}_{X \cup (M \setminus U)}^r(M)$ by right composition.

Corollary 4.1. (1) The image $\pi(\mathcal{D}_{X \cup (M \setminus U)}^r(M))$ is open and closed in $\mathcal{E}_X^r(N, U)$ and the map

$$\pi : \mathcal{D}_{X \cup (M \setminus U)}^r(M) \rightarrow \pi(\mathcal{D}_{X \cup (M \setminus U)}^r(M))$$

is a principal bundle with fiber $\mathcal{D}_{N \cup (M \setminus U)}^r(M)$.

(2) The restriction map

$$\pi : \mathcal{D}_{X \cup (M \setminus U)}^r(M)_0 \rightarrow \mathcal{E}_X^r(N, U)_0, \quad \pi(h) = h|_N,$$

is a principal bundle with fiber $\mathcal{D}_{X \cup (M \setminus U)}^r(M)_0 \cap \mathcal{D}_N(M)$.

4.3. Diffeomorphism groups of noncompact n -manifolds.

Suppose M is a noncompact connected smooth n -manifold without boundary and X is a compact smooth submanifold of M . A *smooth exhausting sequence* of (M, X) means an exhausting sequence $\{M_i\}_{i \geq 1}$ of M such that each M_i is a compact n -submanifold of M for each $i \geq 1$ and $X \subset \text{Int } M_1$. Let $M_0 = X$. Obviously (M, X) has a smooth exhausting sequence and any smooth exhausting sequence of (M, X) satisfies the assumption (A) with respect to the diffeomorphism group $\mathcal{D}_X(M)$. Therefore, Theorem 3.1 and Lemma 4.1 (3) yield the following consequences.

Proposition 4.1. Suppose M admits a smooth exhausting sequence $\{M_i\}_{i \geq 1}$ such that

- (i) $\mathcal{D}_{M_k \cup (M - U_i)}(M)_0 \simeq *$ for each $i > k \geq 0$,
- (ii) $\mathcal{D}_{M_k \cup (M - U_j)}(M)_0 \cap \mathcal{D}_{M_i}(M) = \mathcal{D}_{M_i \cup (M - U_j)}(M)_0$ for each $j > i > k \geq 0$.

Then, (1) $\mathcal{D}_X(M)_0 \cong \ell^2$ and (2) $\mathcal{D}_X(M)_0^{c*}$ is HD in $\mathcal{D}_X(M)_0$.

Proposition 4.2. If $M = \text{Int } N$ for some compact n -manifold N with non-empty boundary, then

- (1) $\mathcal{D}_X(M)_0$ is an ℓ^2 -manifold,
- (2) if $\mathcal{D}_{\partial N \times \{0\}}(\partial N \times [0, 1]) \simeq *$, then $\mathcal{D}_X(M)_0 \cong \mathcal{D}_X(N)_0$,
- (3) if $\mathcal{D}_{\partial N \times \{0, 1\}}(\partial N \times [0, 1])_0 \simeq *$, then $\mathcal{D}_X(M)_0^{c*}$ is HD in $\mathcal{D}_X(M)_0$.

Proof. Take a smooth closed collar $\partial N \times [0, 2]$ of $\partial N = \partial N \times \{0\}$ in $N - X$ and let $M_i = M - (\partial N \times [0, 1/i])$ ($i \geq 1$). Then $\{M_i\}_{i \geq 1}$ is a smooth exhausting sequence of (M, X) . Let $G = \mathcal{D}_X^r(M)$. The tuple $(G, M, \{M_i\}_{i \geq 1})$ satisfies the assumption (A).

(1) By Theorem 3.1 (2)(i) and Lemma 4.1 (3) it suffices to show that $G_{M_i} = \mathcal{D}_{M_i}^r(M) \simeq *$ for each $i \geq 1$. This is verified by the Alexander trick towards ∞ , since $M - \text{Int } M_i = \partial N \times (0, 1/i]$ is an open collar of $\partial M_i = \partial N \times \{1/i\}$.

(2) By (1) and Lemma 4.2, both $\mathcal{D}_X^r(M)_0$ and $\mathcal{D}_X^r(N)_0$ are ℓ^2 -manifolds. Thus, by Theorem 2.1 (2) it suffices to show that $\mathcal{D}_X^r(M)_0 \simeq \mathcal{D}_X^r(N)_0$.

Note that (i) G_{M_1} is an AR and $(G_{M_1})_0 = G_{M_1}$ by (1) and (ii) $\mathcal{D}_{M_1}^r(N)$ is an AR since it is an ANR and $\mathcal{D}_{M_1}^r(N) \simeq \mathcal{D}_{\partial N \times \{1\}}(\partial N \times [0, 1]) \simeq *$ by the assumption. The restriction maps

$$\pi : \mathcal{D}_X^r(M)_0 \longrightarrow \mathcal{E}_X^r(M_1, M)_0 \quad \text{and} \quad \pi_1 : \mathcal{D}_X^r(N)_0 \longrightarrow \mathcal{E}_X^r(M_1, M)_0$$

are principal bundles with the AR fibers

$$G_0 \cap G_{M_1} = G_{M_1} \quad \text{and} \quad \mathcal{D}_X^r(N)_0 \cap \mathcal{D}_{M_1}^r(N) = \mathcal{D}_{M_1}^r(N).$$

Thus, by Lemma 2.2 the maps π and π_1 are homotopy equivalences.

(3) Let $\mathcal{H} = \mathcal{D}^r(\partial N \times \mathbb{R})$. Then, for $j > i \geq 2$ we have

$$\begin{aligned} G_{M_1}(U_j)_0 \cap G_{M_i} &\cong \mathcal{H}(\partial N \times [0, 2])_0 \cap \mathcal{H}(\partial N \times [1, 2]) = \mathcal{H}(\partial N \times [1, 2])_0 \\ &\simeq \mathcal{D}_{\partial N \times \{0, 1\}}(\partial N \times [0, 1])_0 \simeq *. \end{aligned}$$

Since $(G_{M_1})_0 = G_{M_1}$, the assertion follows from Theorem 3.1 (2)(ii). $\square$

Proposition 1.2 now follows from Proposition 4.2 (1). In the next section, Propositions 4.1 and 4.2 are used to deduce Theorems 1.1 and 1.2.

5. DIFFEOMORPHISM GROUPS OF NON-COMPACT 2-MANIFOLDS

5.1. Fundamental facts on diffeomorphism groups of 2-manifolds.

First we recall some fundamental facts on diffeomorphism groups of compact 2-manifolds. The symbols $\mathbb{S}^1$, $\mathbb{S}^2$, $\mathbb{T}$, $\mathbb{D}$, $\mathbb{A}$, $\mathbb{P}$, $\mathbb{K}$ and $\mathbb{M}$ denote the 1-sphere (circle), 2-sphere, torus, disk, annulus, projective plane, Klein bottle and Möbius band, respectively.

Theorem 5.1. ([9, 26] etc.) *Suppose M is a compact connected smooth 2-manifold (possibly with boundary) and X is a compact smooth submanifold of M .*

- (i) (a) $\mathcal{D}^r(M)_0 \simeq SO(3)$ if $M \cong \mathbb{S}^2, \mathbb{P}$. (b) $\mathcal{D}^r(M)_0 \simeq \mathbb{T}$ if $M \cong \mathbb{T}$.
- (ii) $\mathcal{D}_X^r(M)_0 \simeq \mathbb{S}^1$ if $(M, X) \cong (\mathbb{S}^2, 1\text{pt}), (\mathbb{S}^2, 2\text{pt}), (\mathbb{D}, \emptyset), (\mathbb{D}, 0), (\mathbb{A}, \emptyset), (\mathbb{M}, \emptyset), (\mathbb{K}, \emptyset)$.
- (iii) $\mathcal{D}_X^r(M)_0 \simeq *$ in all other cases.
- (iv) $\mathcal{D}_\partial^r(\mathbb{D}) \simeq \mathcal{D}_{\{0\} \cup \partial}^r(\mathbb{D}) \simeq *$ and $\mathcal{D}_\partial^r(\mathbb{M}) \simeq *$.

By [9] and a C^r -analogue of [10] we have

Lemma 5.1. *Suppose M is a compact smooth 2-manifold (possibly with boundary).*

- (1) *Suppose N is a closed collar of ∂M . If $h \in \mathcal{D}_N^r(M)$ is homotopic to id_M rel N , then h is C^r -isotopic to id_M rel N and $h \in \mathcal{D}_N^r(M)_0$.*
- (2) *Suppose N is a compact smooth 2-submanifold of M with $\partial M \subset N$.*
 - (i) *If $h \in \mathcal{D}_N^r(M)$ is C^0 -isotopic to id_M rel N , then h is C^r -isotopic to id_M rel N and $h \in \mathcal{D}_N^r(M)_0$.*
 - (ii) *$h \in \mathcal{D}_N^r(M)_0$ iff $h \in \mathcal{D}_N^r(M)$ and h is C^r -isotopic to id_M rel N (cf. Remark 2.1 (2)).*

In Corollary 4.1 we have a principal bundle with fiber $\mathcal{G} \equiv \mathcal{D}_X^r(M)_0 \cap \mathcal{D}_N^r(M)$. The next theorem provides us with a sufficient condition which implies $\mathcal{G} = \mathcal{D}_N^r(M)_0$. The symbol $\#X$ denotes the number of elements (or cardinal) of a set X .

Theorem 5.2. *Suppose M is a compact connected smooth 2-manifold (possibly with boundary), N is a compact smooth 2-submanifold of M with $\partial M \subset N$ and X is a subset of N . Suppose (M, N, X) satisfies the following conditions:*

- (i) $M \neq \mathbb{T}, \mathbb{P}, \mathbb{K}$ or $X \neq \emptyset$.
- (ii) (a) *if H is a disk component of N , then $\#(H \cap X) \geq 2$,*
(b) *if H is an annulus or Möbius band component of N , then $H \cap X \neq \emptyset$,*

- (iii) (a) if L is a disk component of $cl(M \setminus N)$, then $\#(L \cap X) \geq 2$,
- (b) if L is a Möbius band component of $cl(M \setminus N)$, then $L \cap X \neq \emptyset$.

Then we have

- (1) if $h \in \mathcal{D}_N^r(M)$ is C^0 -isotopic to id_M rel X , then h is C^r -isotopic to id_M rel N ,
- (2) $\mathcal{D}_X^r(M)_0 \cap \mathcal{D}_N^r(M) = \mathcal{D}_N^r(M)_0$.

Lemma 5.2. *Suppose M is a connected smooth 2-manifold without boundary and N is a smooth closed 2-submanifold of M . Suppose $N \neq \emptyset$, $cl(M \setminus N)$ is compact and (M, N) satisfies the following conditions:*

- (i) $M \not\cong \mathbb{T}, \mathbb{P}, \mathbb{K}$,
- (ii) each component C of ∂N does not bound a disk or a Möbius band,
- (iii) each component of $N \not\cong \mathbb{S}^1 \times [0, 1], \mathbb{S}^1 \times [0, 1)$.

If $h \in \mathcal{D}_N^r(M)$ is C^0 -isotopic to id_M , then h is C^r -isotopic to id_M rel N and $h \in \mathcal{D}_N^r(M)_0$.

Theorem 5.2 and Lemma 5.2 follow from Lemma 5.1 and the corresponding statements in the C^0 -case [29, Theorem 3.1, Lemma 3.4 (and a remark after Lemma 3.4)].

5.2. Diffeomorphism groups of noncompact 2-manifolds.

In this section we prove Theorems 1.1 and 1.2. Suppose M is a noncompact connected smooth 2-manifold without boundary and X is a compact smooth submanifold of M (i.e., a disjoint union of a compact smooth 2-submanifold of M and finitely many smooth circles and points). We need to separate the next two cases:

- (I) $(M, X) = (\text{a plane}, \emptyset), (\text{a plane}, 1 \text{ pt}), (\text{an open Möbius band}, \emptyset) \text{ or } (\text{an open annulus}, \emptyset)$.
- (II) (M, X) is not Case (I).

Theorems 1.1 and 1.2 are rewritten as follows:

Theorem 5.3. (1) $\mathcal{D}_X^r(M)_0$ is an ℓ_2 -manifold.

- (2) $\mathcal{D}_X^r(M)_0 \simeq \mathbb{S}^1$ in Case (I) and $\mathcal{D}_X^r(M)_0 \simeq *$ in Case (II).
- (3) $\mathcal{D}_X^r(M)_0^{c*}$ is HD in $\mathcal{D}_X^r(M)_0$.

Proof. Case (I): From the assumption it follows that $(M, X) = (\text{Int } N, X)$, where $(N, X) = (\mathbb{D}, \emptyset), (\mathbb{D}, 1 \text{ pt}), (\mathbb{M}, \emptyset) \text{ or } (\mathbb{A}, \emptyset)$. By Theorem 5.1 it is seen that $\mathcal{D}_X^r(N)_0 \simeq \mathbb{S}^1$ and

$$\mathcal{D}_{L \times \{0\}}^r(L \times [0, 1]) = \mathcal{D}_{L \times \{0\}}^r(L \times [0, 1])_0 \simeq * \quad \text{and} \quad \mathcal{D}_{L \times \{0, 1\}}^r(L \times [0, 1])_0 \simeq *$$

for any closed smooth 1-manifold L (i.e., L is a disjoint union of finitely many circles). Therefore, from Proposition 4.2 for $n = 2$ follows the assertions (1), (3) and (2) (I) $\mathcal{D}_X^r(M)_0 \cong \mathcal{D}_X^r(N)_0 \simeq \mathbb{S}^1$.

Case (II): We can write as $M = \cup_{i=0}^{\infty} M_i$, where $M_0 = X$ and for each $i \geq 1$

- (a) M_i is a nonempty compact connected smooth 2-submanifold of M and $M_{i-1} \subset \text{Int } M_i$,
- (b) for each connected component L of $cl(M \setminus M_i)$, L is noncompact and $L \cap M_{i+1}$ is connected.

Note that M is a plane (an open Möbius band, an open annulus) iff infinitely many M_i 's are disks (Möbius bands, annuli respectively), and that any subsequence of M_i ($i \geq 1$) also satisfies the conditions (a) and (b). Thus, passing to a subsequence, we may assume that

- (i) if M is a plane, then each M_i is a disk,
- (ii) if M is an open Möbius band, then each M_i is a Möbius band,
- (iii) if M is an open annulus, then each M_i is an annulus (and the inclusion $M_i \subset M_{i+1}$ is essential),
- (iv) if M is not a plane, an open Möbius band or an open annulus, then each M_i is not a disk, an annulus or a Möbius band.

For each $i \geq 1$ let $U_i = \text{int } M_i$.

To apply Proposition 4.1 we have to verify the following conditions:

- (b)₁ $\mathcal{D}_{M_k \cup (M - U_i)}^r(M)_0 \simeq *$ for each $i > k \geq 0$,
- (b)₂ $\mathcal{G}_{k,j}^i \equiv \mathcal{D}_{M_k \cup (M - U_j)}^r(M)_0 \cap \mathcal{D}_{M_i}^r(M) = \mathcal{D}_{M_i \cup (M - U_j)}^r(M)_0$ for each $j > i > k \geq 0$.

Choose a small closed collar E_i of ∂M_i in $U_{i+1} \setminus U_i$ and set $M'_i = M_i \cup E_i \subset U_{i+1}$.

- (b)₁: $\mathcal{D}_{M_k \cup (M - U_i)}^r(M)_0 \cong \mathcal{D}_{M_k \cup E_i}^r(M'_i) \simeq *$ by Theorem 5.1.
- (b)₂: It suffices to show that

$$\mathcal{D}_{M_k \cup E_j}^r(M'_j)_0 \cap \mathcal{D}_{M_i \cup E_j}^r(M'_j) = \mathcal{D}_{M_i \cup E_j}^r(M'_j)_0.$$

We apply Theorem 5.2 to $(\tilde{M}, \tilde{N}, \tilde{X}) = (M'_j, M_i \cup E_j, M_k \cup E_j)$. It remains to verify that this triple satisfies the conditions (i)–(iii) in Theorem 5.2:

(ii): The components of $\tilde{N}$ consists of M_i and the annulus components of E_j . The latter components obviously satisfy the condition (ii) (b). By the choice of $\{M_j\}_{j \geq 1}$, if $M_i \cong \mathbb{D} (\mathbb{M}, \mathbb{A})$, then M is a plane (an open Möbius band, an open annulus). Thus the assumption of Case (II) implies that $(M_i, X) \neq (\mathbb{D}, \emptyset), (\mathbb{D}, 1pt), (\mathbb{M}, \emptyset), (\mathbb{A}, \emptyset)$ and that M_i satisfies the condition (ii).

(iii): Each component L of $cl(\tilde{M} \setminus \tilde{N}) = M_j \setminus U_i$ meets both ∂M_i and ∂M_j . Thus ∂L is not connected and $L \not\cong \mathbb{D}, \mathbb{M}$. This completes the proof. $\square$

Proof of Corollary 1.2. Since $\mathcal{D}_X^r(M)_0^{c*} \subset \mathcal{D}_X^r(M)_0^c \subset \mathcal{D}_X^r(M)_0$, the subgroup $\mathcal{D}_X^r(M)_0^c$ is also HD in $\mathcal{D}_X^r(M)_0$. Thus, the assertion (1) follows from Theorem 1.1 (1) and Lemma 2.3 (1)(ii), while the assertion (2) follows from Lemma 2.3 (1)(i). $\square$

Proof of Proposition 1.1. For notational simplicity we set $G = \mathcal{D}_X^r(M)$. We choose an exhausting sequence $\{M_i\}_{i \geq 1}$ of (M, X) as in the proof of Theorem 5.3 Case (II).

[A] If (M, X) satisfies the condition (a) or (b), then $G_0^c = G_0^{c*}$:

Given $h \in G_0^c$, there exists $i \geq 1$ such that $h \in G(U_i)$. We show that $h \in G(U_i)_0 \subset G_0^{c*}$. If (M, X) satisfies the condition (b), then by Theorem 5.1 (iv) $G(U_i) \cong \mathcal{D}_{E_i}^r(M_i \cup E_i) \simeq \mathcal{D}_{\partial M_i}^r(M_i) \simeq *$, where E_i is a small closed collar of ∂M_i in $M \setminus U_i$. This means that $G(U_i) = G(U_i)_0$. Below we assume that (M, X) does not satisfy the condition (b). Note that h is C^0 -isotopic to $id_M \text{ rel } X$ since $h \in G_0$ and G_0 is a connected ANR.

(1) The case (a) with $X = \emptyset$.

Let $N = M \setminus U_i$ and we apply Lemma 5.2 to (M, N) and h . The conditions (ii) and (iii) in Lemma 5.2 are verified as follows:

(ii) If a circle component C of $\partial N = \partial M_i$ bounds a disk or a Möbius band D , then $M_i = D$ and by the choice of the exhausting sequence $\{M_i\}_{i \geq 1}$ it follows that M is a plane or an open Möbius band. This contradicts the assumption that (M, X) does not satisfy the condition (b)

(iii) Any component of N is noncompact and not diffeomorphic to $\mathbb{S}^1 \times [0, 1)$ by the condition (a).

Since $h \in G_0$ is C^0 -isotopic to id_M , from Lemma 5.2 it follows that h is C^r -isotopic to id_M rel N and $h \in G(U_i)_0$.

(2) The case (a) with $X \neq \emptyset$:

Let F_k ($k = 1, \dots, m$) denote the connected components of the compact 2-submanifold $cl(M_i \setminus X)$ which are disks or Möbius bands let H_ℓ ($\ell = 1, \dots, n$) denote the remaining components. Set $N = X \cup (\cup_k F_k) \cup (M \setminus U_i)$. There exists a C^0 -isotopy h_t rel X from h to id_M . Since $\partial F_k \subset X$, it follows that h_t maps each F_k onto itself. Thus, if we define $h' \in G_N$ by $h' = h$ on $X \cup (\cup_\ell H_\ell) \cup (M \setminus U_i)$ and $h' = id$ on $\cup_k F_k$, then h' is C^0 -isotopic to h rel $X \cup (M \setminus U_i)$. and C^0 -isotopic to id_M rel X .

Define $A \subset X$ by choosing two interior points from each component of X . Set $M' = M \setminus A$ and $N' = N \setminus A$. Then $h'|_{M'} \in \mathcal{D}_{N'}^r(M')$ is C^0 -isotopic to $id_{M'}$. We show that (M', N') satisfies the conditions (i)–(iii) in Lemma 5.2. First note that M' is noncompact and $L \equiv cl(M' \setminus N') = cl(M \setminus N) = \cup_\ell H_\ell$ is compact.

(ii) Suppose a circle component C of $\partial N = \partial M_i \cup (\partial X \setminus (\cup_k \partial F_k))$ bounds a disk or a Möbius band D in M' . Then M is the union of D and a connected submanifold $W = cl(M - D)$ with $\partial W = C$, and it follows that $N \subset W$. In fact, (α) $X \subset W$ since $A \subset W$ and each component of X meets A , (β) each $F_k \subset W$ since F_k meets X but does not meet C and (γ) $M \setminus U_i \subset W$ since each component of $M \setminus U_i$ is noncompact. If $C \subset \partial X \setminus (\cup_k \partial F_k)$, Thus $D \subset \cup_\ell H_\ell$ and $D = H_\ell$ for some ℓ , but this contradicts the choice of H_ℓ . If $C \subset \partial M_i$, then $D \supset M_i \supset A$, which contradicts that $D \subset M'$.

(iii) Let J be any component of N' . If $J \subset (X \setminus A) \cup (\cup_k F_k)$, then J has two ends, since it is the union of $X_1 \setminus A$ for some component X_1 of X and some F_k 's. Otherwise, J is a component of $M \setminus U_i$, so it is noncompact and $J \not\cong \mathbb{S}^1 \times [0, 1)$ by the condition (a).

Therefore, by Lemma 5.2 $h'|_{M'}$ is C^0 -isotopic to $id_{M'}$ rel N' and by the end compactification we obtain a C^0 -isotopy $h \simeq h' \simeq id_M$ rel $X \cup (M \setminus U_i)$. This implies that $h \in G(U_i)_0$.

[B] If (M, X) does not satisfy the conditions (a) and (b), then $G_0^{c*} \subsetneq G_0^c$:

Suppose (M, X) does not satisfy the conditions (a) and (b). Then M contains a product end $E \cong \mathbb{S}^1 \times [0, 1)$ such that $E \cap X = \emptyset$. We identify E with $\mathbb{S}^1 \times [0, 1)$. Note that $F = cl(M \setminus E)$ is a connected submanifold of M and $\partial F = \partial E$. Let $h \in G^c$ denote a Dehn twist in E . We show that $h \in G_0^c \setminus G_0^{c*}$. A C^r -isotopy $h_t : h \simeq id_M$ with $h_t \in G_0^c$ is obtained by sliding the Dehn twist towards ∞ . This implies that $h \in G_0^c$. It remains to show that $h \notin G_0^{c*}$. On the contrary, suppose $h \in G_0^{c*}$. Then $h \in G(M_i)_0$ for some $i \geq 1$ and there exists a C^0 -isotopy $h_t : h \simeq id_M$ rel X with $\text{supp } h_t \subset M_i$ ($0 \leq t \leq 1$).

If $F \cong \mathbb{S}^1 \times [0, 1)$, then $M \cong \mathbb{S}^1 \times (-1, 1)$ and M_i is contained in an annulus L in M . This means that the Dehn twist in L is C^0 -isotopic to id_L rel ∂L , a contradiction. Thus we may assume that $F \not\cong \mathbb{S}^1 \times [0, 1)$.

Choose $t \in (0, 1)$ such that $M_i \subset F \cup L$, where $L = \mathbb{S}^1 \times [0, t]$ is an annulus. Let $Y = \mathbb{S}^1 \times [t, 1)$ ($= cl(E \setminus L)$) and $N = F \cup Y$. Then $h|_N = id$ and $h_t : h \simeq id_M$ is a C^0 -isotopy rel $X \cup Y$. Define A by choosing an interior point from Y and two interior points from X if $X \neq \emptyset$. Let $M' = M \setminus A$ and $N' = N \setminus A$. We show that (M', N') satisfies the conditions (i)–(iii) in Lemma 5.2.

(ii) $\partial N' = \partial N$ consists of two circles $C_1 = \mathbb{S}^1 \times \{0\}$ and $C_2 = \mathbb{S}^1 \times \{t\}$. If C_1 bounds a disk or a Möbius band D in M' , then $F = D \subset M'$, so we have $X = \emptyset$ and $M = D \cup E$ is a plane or an open Möbius band (i.e. the condition (b)). This contradicts the assumption in [B]. Similarly, the circle C_2 bound neither a disk nor a Möbius band in M' .

(iii) Note that N' consists of the two components $F \setminus A$ and $Y \setminus A$. These are homeomorphic to neither $\mathbb{S}^1 \times [0, 1]$ nor $\mathbb{S}^1 \times [0, 1)$. In fact, $Y \setminus A$ has two ends, and if $X \neq \emptyset$, then $F \setminus A$ has at least two ends. If $X = \emptyset$, then $F \setminus A = F \not\simeq \mathbb{S}^1 \times [0, 1)$ and ∂F is a circle.

Since $h' = h|_{M'}$ is C^0 -isotopic to $id_{M'}$ and $h'|_{N'} = id$, by Lemma 5.2 we obtain a C^0 -isotopy $h' \simeq id_{M'}$ rel N' . This means that the Dehn twist $h|_L$ on the annulus L is isotopic to id_L rel ∂L . This is a contradiction. This completes the proof. $\square$

6. GROUPS OF VOLUME-PRESERVING DIFFEOMORPHISMS OF NONCOMPACT 2-MANIFOLDS

In this final section we discuss topological types of groups of volume-preserving diffeomorphisms of noncompact 2-manifolds with the compact-open C^∞ -topology. Since we are only concerned with the groups of C^∞ -diffeomorphisms with the compact-open C^∞ -topology, we omit the symbol ∞ from the notations.

6.1. General properties of groups of volume-preserving diffeomorphisms of n -manifolds.

Suppose M is a connected oriented smooth n -manifold possibly with boundary, $X \subsetneq M$ is a closed subset of M and ω is a positive volume form on M . Let $\mathcal{D}_X(M; \omega) \subset \mathcal{D}_X^+(M)$ denote the subgroups of $\mathcal{D}_X(M)$ consisting of ω -preserving diffeomorphisms and orientation-preserving diffeomorphisms of M respectively, that is,

$$\mathcal{D}_X(M; \omega) = \{h \in \mathcal{D}_X(M) \mid h^*\omega = \omega\}.$$

These subgroups are endowed with the subspace topology (i.e., the compact-open C^∞ -topology). As before, the subscript ‘0’ denotes the identity connected component. Note that $\mathcal{D}_X^+(M)_0 = \mathcal{D}_X(M)_0$. The group $\mathcal{D}_X(M; \omega)$ is a separable, completely metrizable topological group since it is a closed subgroup of $\mathcal{D}_X(M)$. It is also seen to be infinite-dimensional and non locally compact. Hence, by Theorem 2.1 the group $\mathcal{D}_X(M; \omega)$ (or $\mathcal{D}_X(M; \omega)_0$) is an ℓ_2 -manifold iff it is an ANR.

First we recall Moser’s theorem [21] and its extension to the noncompact case [31]. (We refer to [1, 5, 31] for end compactifications and related matters.) The space E_M of ends of M is a compact 0-dimensional metrizable space. Let E_M^ω denote the subspace of E_M consisting of ω -finite ends of M . Each $h \in \mathcal{H}(M)$ admits a unique homeomorphic extension $\bar{h}$ on the end compactification $M \cup E_M$. We define $\mathcal{D}^+(M, E_M^\omega) = \{h \in \mathcal{D}^+(M) \mid \bar{h}(E_M^\omega) = E_M^\omega\}$.

Let $\mathcal{V}^+(M; \omega(M), E_M^\omega)_{ew}$ denote the space of positive volume forms μ on M such that $\mu(M) = \omega(M)$ and $E_M^\mu = E_M^\omega$. This space is endowed with the finite-ends weak C^∞ -topology ew (cf. [5, 31]).

The group $\mathcal{D}^+(M, E_M^\omega)$ acts on the space $\mathcal{V}^+(M; \omega(M), E_M^\omega)$ by the push-forward of forms and induces the orbit map at ω ,

$$\pi_\omega : \mathcal{D}^+(M, E_M^\omega) \longrightarrow \mathcal{V}^+(M; \omega(M), E_M^\omega)_{ew}.$$

Moser's theorem [21] and [31, Corollary 1.1] assert that this orbit map admits a section into $\mathcal{D}_\partial(M)_0$. This implies the following relation between the groups $\mathcal{D}_X(M; \omega) \subset \mathcal{D}_X(M)$.

Proposition 6.1.

- (1) $(\mathcal{D}^{\infty+}(M; E_M^\omega), \mathcal{D}(M; \omega)) \cong (\mathcal{V}^+(M; \omega(M), E_M^\omega)_{ew}, \{\omega\}) \times \mathcal{D}(M; \omega)$,
 $(\mathcal{D}(M)_0, \mathcal{D}(M; \omega)_0) \cong (\mathcal{V}^+(M; \omega(M), E_M^\omega)_{ew}, \{\omega\}) \times \mathcal{D}(M; \omega)_0$.
- (2) Suppose N is a compact n -submanifold of $\text{Int } M$.
 - (i) $\mathcal{D}_{M-N}(M; \omega)$ is a SDR of $\mathcal{D}_{M-N}^+(M)$ and is an ANR (since $\mathcal{D}_{M-N}(M)$ is an ANR).
 - (ii) $\mathcal{D}_N(M; \omega)$ is a SDR of $\mathcal{D}_N^+(M, E_M^\omega)$ and $\mathcal{D}_N(M; \omega)_0$ is a SDR of $\mathcal{D}_N(M)_0$.

In the statement (2) we apply Moser's theorem [21] and [31, Corollary 1.1] to each $L \in \mathcal{C}(M - \text{Int } N)$.

Next we recall the definition of the end charge homomorphism introduced by S. R. Alpern and V.S. Prasad [1]. Suppose M is a noncompact connected orientable smooth n -manifold possibly with boundary and ω is a volume form on M . An end charge of M is a finitely additive signed measure on the algebra of clopen subsets of E_M . Let $\mathcal{S}(M)$ denote the topological linear space of all end charges of M with the weak topology and let $\mathcal{S}(M; \omega)$ denote the linear subspace of $\mathcal{S}(M)$ consisting of end charges c of M with $c(E_M) = 0$ and $c|_{E_M^\omega} = 0$.

Let $\mathcal{D}_{E_M}(M; \omega) = \{h \in \mathcal{D}(M; \omega) \mid \bar{h}|_{E_M} = \text{id}_{E_M}\}$. We have $\mathcal{D}_{E_M}(M; \omega)_0 = \mathcal{D}(M; \omega)_0$. For each $h \in \mathcal{D}_{E_M}(M; \omega)$ an end charge $c^\omega(h) \in \mathcal{S}(M; \omega)$ is defined by

$$c^\omega(h)(E_C) = \omega(C - h(C)) - \omega(h(C) - C),$$

where C is any n -submanifold of M such that $\text{Fr}_M C$ is compact and $E_C \subset E_M$ is the set of ends of C . The end charge homomorphism

$$c^\omega : \mathcal{D}_{E_M}(M; \omega) \rightarrow \mathcal{S}(M; \omega) : h \longmapsto c^\omega(h)$$

is a continuous group homomorphism. Let $c_0^\omega : \mathcal{D}(M; \omega)_0 \rightarrow \mathcal{S}(M; \omega)$ denote the restriction of c^ω to $\mathcal{D}(M; \omega)_0$.

The kernels $\ker c^\omega$ and $\ker c_0^\omega$ are separable, non locally compact, completely metrizable topological groups. Hence, by Theorem 2.1 the group $\ker c^\omega$ (or $\ker c_0^\omega$) is an ℓ_2 -manifold iff it is an ANR. In [31, Corollary 1.2] we have shown that the homomorphism c^ω has a continuous (non-homomorphic) section into $\mathcal{D}_\partial(M; \omega)_0$. This clarifies the relation between the groups $\ker c_0^\omega \subset \mathcal{D}(M; \omega)_0$.

Proposition 6.2.

- (1) $(\mathcal{D}_{E_M}(M; \omega), \ker c^\omega) \cong (\mathcal{S}(M; \omega), 0) \times \ker c^\omega$, $(\mathcal{D}(M; \omega)_0, \ker c_0^\omega) \cong (\mathcal{S}(M; \omega), 0) \times \ker c_0^\omega$.
- (2) (i) $\ker c^\omega$ is a SDR of $\mathcal{D}_{E_M}(M; \omega)$.
(ii) $\ker c_0^\omega = (\ker c^\omega)_0$ and it is a SDR of $\mathcal{D}(M; \omega)_0$.

6.2. The bundle theorem.

Next we obtain the bundle theorems for groups of volume-preserving diffeomorphisms. Suppose M is a connected oriented smooth n -manifold without boundary, ω is a positive volume form on M , N is a compact smooth n -submanifold of M , X is a closed subset of N and N_0 is smooth n -submanifold of M such that $N \subset U_0 \equiv \text{Int } N_0$.

For notational simplicity we set $(G, H, F) = (\mathcal{D}(M), \mathcal{D}(M; \omega), \ker c^\omega)$. Let $\mathcal{V}^+(M)$ denote the space of positive volume forms on M endowed with the compact-open C^∞ -topology. The group G_0 acts continuously on $\mathcal{V}^+(M)$ from the right by the pullback $\mu \cdot h = h^*\mu$. For any subset $\mathcal{F}$ of $\mathcal{E}_X^\infty(N, M)$ the symbol $\mathcal{F}^{co}$ denotes the space $\mathcal{F}$ endowed with the subspace topology (= the compact-open C^∞ -topology). When $i_N \in \mathcal{F}^{co}$, the symbol $\mathcal{F}_0^{co}$ denotes the connected component of the inclusion $i_N : N \subset M$ in $\mathcal{F}^{co}$. Hence, the space $\mathcal{E}_X^H(N, U_0)$ carries the quotient topology, while $\mathcal{E}_X^H(N, U_0)^{co}$ carries the compact-open C^∞ -topology. Let $\mathcal{C}(Y)$ denote the set of connected components of a space Y .

The extension theorems for the transformation groups H and F are summarized as follows:

Theorem 6.1. (H) *There exists a neighborhood $\mathcal{U}$ of i_N in $\mathcal{E}_X^H(N, U_0)^{co}$ and*

a map $\varphi : \mathcal{U} \rightarrow H(U_0)_0 \cap H_X$ such that $\varphi(f)|_N = f$ ($f \in \mathcal{U}$) and $\varphi(i_N) = id_M$.

(F) *Suppose ∂N_0 is compact and U is an open neighborhood of N in U_0 such that $U \cap L$ is connected for each $L \in \mathcal{C}(N_0 - \text{Int } N)$. Then there exists a neighborhood $\mathcal{U}$ of i_N in $\mathcal{E}_X^F(N, U_0)^{co}$ and a map $\varphi : \mathcal{U} \rightarrow F(U)_0 \cap F_X$ such that $\varphi(f)|_N = f$ ($f \in \mathcal{U}$) and $\varphi(i_N) = id_M$.*

Proof. In each case we may assume that $X = \emptyset$, since $\varphi(f)|_X = i_X$ if $\varphi(f)|_N = f$ and $f|_X = i_X$.

(H) Choose any compact smooth n -submanifold N_1 of U_0 with $N \subset U_1 \equiv \text{Int } N_1$. Consider the subspace of $\mathcal{V}^+(M)$ defined by

$$\mathcal{V}_N^\omega(U_1) = \{\mu \in \mathcal{V}^+(M) \mid \mu = \omega \text{ on } N \cup (M - U_1) \text{ and } \mu(L) = \omega(L) \text{ for each } L \in \mathcal{C}(N_0 - \text{Int } N)\}.$$

Applying Moser's theorem [21] or [31, Corollary 1.1] to each $L \in \mathcal{C}(N_0 - \text{Int } N)$ we obtain a map

$$\eta : \mathcal{V}_N^\omega(U_1) \rightarrow G_N(U_0)_0$$

such that $\eta(\mu)^*\mu = \omega$ ($\mu \in \mathcal{V}_N^\omega(U_1)$) and $\eta(\omega) = id_M$.

By Theorem 4.1 there exists a neighborhood $\mathcal{U}_1$ of i_N in $\mathcal{E}^\infty(N, U_1)$ and a map

$$\psi : \mathcal{U}_1 \rightarrow G(U_1)_0$$

such that $\psi(f)|_N = f$ ($f \in \mathcal{U}_1$) and $\psi(i_N) = id_M$. Then $\mathcal{U} = \mathcal{U}_1 \cap \mathcal{E}^H(N, U_0)^{co}$ is a neighborhood of i_N in $\mathcal{E}^H(N, U_0)^{co}$.

The map ψ induces a map

$$\chi : \mathcal{U} \rightarrow \mathcal{V}_N^\omega(U_1), \quad \chi(f) = \psi(f)^*\omega.$$

We verify that $\chi(f) \in \mathcal{V}_N^\omega(U_1)$ ($f \in \mathcal{U}$). Since $f \in \mathcal{E}^H(N, U_0)$ there exists a $h \in H(U_0)$ with $h|_N = f$. Since $\psi(f)|_N = f$ and $\psi(f) = id$ on $M - U_1$, we have $\chi(f)|_N = f^*\omega = (h^*\omega)|_N = \omega|_N$ and $\chi(f)|_{M-U_1} = \omega|_{M-U_1}$. Since $h^{-1}\psi(f) \in G_N(U_0)$, for each $L \in \mathcal{C}(N_0 - \text{Int } N)$ it follows that $h^{-1}\psi(f)(L) = L$ and $\psi(f)(L) = h(L)$, and that $\chi(f)(L) = \omega(\psi(f)(L)) = \omega(h(L)) = \omega(L)$.

Finally, the required map $\varphi : \mathcal{U} \rightarrow H(U_0)_0$ is defined by

$$\varphi(f) = \psi(f)\eta(\chi(f)) \quad (f \in \mathcal{U}).$$

By Proposition 6.1 (2)(ii) we have $\varphi(f) \in H(U_0) \cap G(U_0)_0 = H(U_0)_0$.

(F) By the assumption we can find a compact smooth n -submanifold N_1 of M such that $N \subset \text{Int } N_1$, $N_1 \subset U$ and $N_1 \cap L$ is connected for each $L \in \mathcal{C}(N_0 - \text{Int } N)$. Let $U_1 = \text{Int } N_1$ and $N_1^* = N_1 - \text{Int } N$.

Consider the subspace of $\mathcal{V}^+(M)$ defined by

$$\mathcal{V}_N^\omega(U_1) = \{\mu \in \mathcal{V}^+(M) \mid \mu = \omega \text{ on } N \cup (M - U_1) \text{ and } \mu(K) = \omega(K) \text{ for each } K \in \mathcal{C}(N_1^*)\}.$$

Applying Moser's theorem [21] to each $K \in \mathcal{C}(N_1^*)$, we obtain a map

$$\eta : \mathcal{V}_N^\omega(U_1) \rightarrow G_N(U_1)_0$$

such that $\eta(\mu)^*\mu = \omega$ ($\mu \in \mathcal{V}$) and $\eta(\omega) = id_M$.

By Theorem 4.1 there exists a neighborhood $\mathcal{U}_1$ of i_N in $\mathcal{E}^\infty(N, U_1)$ and a map

$$\psi : \mathcal{U}_1 \rightarrow G(U_1)_0$$

such that $\psi(f)|_N = f$ ($f \in \mathcal{U}_1$) and $\psi(i_N) = id_M$. Then $\mathcal{U} = \mathcal{U}_1 \cap \mathcal{E}^F(N, U_0)^{co}$ is a neighborhood of i_N in $\mathcal{E}^F(N, M)^{co}$.

The map ψ induces a map

$$\chi : \mathcal{U} \rightarrow \mathcal{V}_N^\omega(U_1), \quad \chi(f) = \psi(f)^*\omega.$$

We show that $\chi(f) \in \mathcal{V}_N^\omega(U_1)$ ($f \in \mathcal{U}$). Since $f \in \mathcal{E}^F(N, U_0)$ there exists an $h \in F(U_0)$ with $h|_N = f$. Since $\psi(f)|_N = f$ and $\psi(f) = id$ on $M - U_1$, we have $\chi(f)|_N = f^*\omega = (h^*\omega)|_N = \omega|_N$ and $\chi(f)|_{M-U_1} = \omega|_{M-U_1}$. By the choice of N_1 , for each $K \in \mathcal{C}(N_1^*)$ there exists a unique $L \in \mathcal{C}(N_0 - \text{Int } N)$ with $K = N_1 \cap L$. Since $h^{-1}\psi(f) \in G_N(U_0)$, it follows that $h^{-1}\psi(f)(L) = L$ and $\psi(f)(L) = h(L)$. Since L is an n -submanifold of M and $\text{Fr } L = \partial L \subset \partial N \cup \partial N_0$ is compact, from the definition of the end charge $c_0^\omega(h)$ it follows that

$$c_0^\omega(h)(E_L) = \omega(L \setminus h(L)) - \omega(h(L) \setminus L).$$

Since $c_0^\omega(h) = 0$, we have $\omega(L \setminus h(L)) = \omega(h(L) \setminus L)$. Let $L_1 = L - N_1$. Then, $L_1 = \psi(f)(L_1) \subset h(L)$ and it is seen that

$$K = L - L_1 = [L \setminus h(L)] \cup [(h(L) \cap L) \setminus L_1] \quad \text{and}$$

$$\psi(f)(K) = \psi(f)(L - L_1) = \psi(f)(L) - L_1 = h(L) - L_1 = [h(L) \setminus L] \cup [(h(L) \cap L) \setminus L_1].$$

Thus we have $\chi(f)(K) = \omega(\psi(f)(K)) = \omega(K)$ and this means that $\chi(f) \in \mathcal{V}_N^\omega(U_1)$.

Since $H(U_1)$ is a SDR of $G(U_1)$, we have $H(U_1) \cap G(U_1)_0 = H(U_1)_0 \subset F(U)_0$. Therefore, the required map $\varphi : \mathcal{U} \rightarrow F(U)_0$ is defined by

$$\varphi(f) = \psi(f)\eta(\chi(f)) \quad (f \in \mathcal{U}).$$

□

Remark 6.1. The statement (F) does not necessarily hold when ∂N_0 is non compact. An example is easily obtained by inspecting a case where two ends of N_0 is included in an end of M .

Consider the restriction map $\pi : F_X(U) \rightarrow \mathcal{E}_X^F(N, U)^{co}$, $\pi(h) = h|_N$. The group $F_N(U)$ acts on $F_X(U)$ by the right translation. The next corollary easily follows from Theorem 6.1 (F) and Lemma 3.1.

Corollary 6.1. *Suppose ∂N_0 is compact and U is an open neighborhood of N in U_0 such that $U \cap L$ is connected for each $L \in \mathcal{C}(N_0 - \text{Int } N)$. Then, the following hold.*

- (1) *The subspace $\mathcal{E}_X^F(N, U)^{co}$ is open in $\mathcal{E}_X^F(N, U_0)^{co}$ and the restriction map*

$$\pi : F_X(U) \rightarrow \mathcal{E}_X^F(N, U)^{co}$$

is a principal bundle with fiber $F_N(U)$.

- (2) *If $F_X(U)_0$ is open in $F_X(U)$, then the subspace $\mathcal{E}_X^F(N, U)_0^{co}$ is closed and open in $\mathcal{E}_X^F(N, U)^{co}$ and the restriction map*

$$\pi : F_X(U)_0 \rightarrow \mathcal{E}_X^F(N, U)_0^{co}$$

is a principal bundle with fiber $F_X(U)_0 \cap F_N$.

We also need the following complementary results to Theorem 6.1.

Lemma 6.1. *Suppose ∂N_0 is compact and $F_X(N_1)_0$ is open in $F_X(N_1)$ for any compact n -submanifold N_1 of U_0 with $X \subset \text{Int } N_1$.*

- (1) *Suppose $f \in \mathcal{E}_X^F(N, U_0)^{co}$ and U is an open neighborhood of $f(N)$ in U_0 such that $U \cap L$ is connected for each $L \in \mathcal{C}(N_0 - \text{Int } f(N))$. Then there exists a neighborhood $\mathcal{V}$ of f in $\mathcal{E}_X^F(N, U_0)^{co}$ and a map $\psi : \mathcal{V} \rightarrow F_X(U)_0$ such that $\psi(g)f = g$ ($g \in \mathcal{V}$) and $\psi(f) = id_M$.*
- (2) *Each $f \in \mathcal{E}_X^F(N, U_0)_0^{co}$ satisfies the following condition.*
- (*) *There exists a compact n -submanifold N_1 of U_0 such that $\mathcal{E}_X^F(N, N_1)_0^{co}$ is a neighborhood of f in $\mathcal{E}_X^F(N, U_0)_0^{co}$.*

Proof. (1) We may assume that $U = \text{Int } N_1$ for some compact n -submanifold N_1 of U_0 . Applying Theorem 6.1 (F) to $(M, N_0, U, f(N), X)$, we obtain a neighborhood $\mathcal{U}$ of $i_{f(N)}$ in $\mathcal{E}_X^F(f(N), U_0)^{co}$ and a map $\varphi : \mathcal{U} \rightarrow F(U)_0 \cap F_X$ such that $\varphi(k)|_{f(N)} = k$ ($k \in \mathcal{U}$) and $\varphi(i_{f(N)}) = id_M$. By the assumption $F_X(U)_0$ is open in $F_X(U)$ and so we may assume that $\varphi(\mathcal{U}) \subset F_X(U)_0$.

Consider the homeomorphism

$$\chi : \mathcal{E}_X^F(f(N), U_0)^{co} \approx \mathcal{E}_X^F(N, U_0)^{co}, \quad \chi(k) = kf.$$

Then $\mathcal{V} = \chi(\mathcal{U})$ and $\psi = \varphi\chi^{-1} : \mathcal{V} \rightarrow F_X(U)_0$ satisfy the required conditions.

(2) We have to show that the subset $\mathcal{F} = \{f \in \mathcal{E}_X^F(N, U_0)_0^{co} \mid (*)\}$ coincides with $\mathcal{E}_X^F(N, U_0)_0^{co}$. Corollary 6.1 implies that $i_N \in \mathcal{F}$. Therefore, it suffices to show that $\mathcal{F}$ is closed and open in $\mathcal{E}_X^F(N, N_1)_0^{co}$. From the definition itself we see that $\mathcal{F}$ is open in $\mathcal{E}_X^F(N, N_1)_0^{co}$.

To see that $\mathcal{F}$ is closed, take any $f \in cl \mathcal{F}$. There exists a compact n -submanifold N_f of U_0 such that $f(N) \subset U_f \equiv \text{Int } N_f$ and $U_f \cap L$ is connected for each $L \in \mathcal{C}(N_0 - \text{Int } f(N))$. By (1) we obtain a neighborhood $\mathcal{V}$ of f in $\mathcal{E}_X^F(N, U_0)_0^{co}$ and a map $\psi : \mathcal{V} \rightarrow F_X(U_f)_0$ such that $\psi(g)f = g$ ($g \in \mathcal{V}$) and $\psi(f) = id_M$. Since $\mathcal{V} \cap \mathcal{F} \neq \emptyset$, we can choose a $g \in \mathcal{V} \cap \mathcal{F}$. Since $g \in \mathcal{V}$ it follows that $\psi(g)f = g$ and $f = \psi(g)^{-1}g$. In turn, since $g \in \mathcal{F}$, there exists a compact n -submanifold N_g of U_0

such that $N \subset U_g \equiv \text{Int } N_g$ and $g \in \mathcal{E}_X^F(N, N_g)_0^{co}$. We may assume that $U_g \cap L$ is connected for each $L \in \mathcal{C}(N_0 - \text{Int } N)$. Then, by Corollary 6.1 the restriction map

$$\pi : F_X(U_g)_0 \rightarrow \mathcal{E}_X^F(N, U_g)_0^{co}$$

is a principal bundle. Hence, there exists an $h \in F_X(U_g)_0$ such that $g = h|_N$.

Take a compact n -submanifold N_1 of U_0 such that $N_f \cup N_g \subset N_1$. Then, we have $\mathcal{V} \subset \mathcal{E}_X^F(N, N_1)_0^{co}$. In fact, for any $k \in \mathcal{V}$, it follows that $\psi(k)\psi(g)^{-1}h \in (F_X(U_f)_0)^2 F_X(U_g)_0 \subset F_X(N_1)_0$ and that $k = \psi(k)f = \psi(k)\psi(g)^{-1}h|_N \in \mathcal{E}_X^F(N, N_1)_0^{co}$. This means that $f \in \mathcal{F}$. This completes the proof. $\square$

6.3. Groups of volume-preserving diffeomorphisms of noncompact n -manifolds.

Suppose M is a noncompact connected orientable smooth n -manifold without boundary, ω is a volume form on M and X is a compact smooth n -submanifold of M . We set $(G, H, F) = (\mathcal{D}(M), \mathcal{D}(M; \omega), \ker c^\omega)$. Choose a smooth exhausting sequence $\{M_i\}_{i \geq 0}$ of M such that $M_0 = X$ and for each $i \geq 1$

- (a) M_i is connected,
- (b) L is noncompact and $L \cap M_{i+1}$ is connected for each $L \in \mathcal{C}(M - \text{Int } M_i)$.

Let $U_i = \text{Int } M_i$ ($i \geq 1$).

Lemma 6.2. *The tuple $(F_X, M, \{M_i\}_{i \geq 1})$ satisfies the assumption (A).*

Proof. (A-0) Since (G, M) has a weak topology, so does (F, M) .

(A-1) Corollary 6.1 implies the following conclusions. For each $j > i > k \geq 0$

- (a) $\mathcal{E}_{M_k}^F(M_i, U_j)^{co} = \mathcal{E}_{M_k}^F(M_i, U_j)$ and $\mathcal{E}_{M_k}^F(M_i, M)^{co} = \mathcal{E}_{M_k}^F(M_i, M)$, since the restriction maps $\pi_{k,j}^i : F_{M_k}(U_j) \rightarrow \mathcal{E}_{M_k}^F(M_i, U_j)^{co}$ and $\pi_k^i : F_{M_k} \rightarrow \mathcal{E}_{M_k}^F(M_i, M)^{co}$ are principal bundles,
- (b) the restriction map $\pi_{k,j}^i : F_{M_k}(U_j)_0 \rightarrow \mathcal{E}_{M_k}^F(M_i, U_j)_0$ is a principal bundle with the structure group $F_{M_k}(U_j)_0 \cap F_{M_i}$.

(A-2) (i) $F_X(U_i) = H_X(U_i)$ is an ANR for each $i \geq 1$.

(A-2)(ii) By (A-1)(a) we can work under the compact-open C^∞ -topology. Let $\mathcal{U}_{k,j}^i = \mathcal{E}_{M_k}^F(M_i, U_j)_0^{co}$ ($j > i > k \geq 0$).

- (a) $\mathcal{U}_{k,j}^i$ is an open subspace of $\mathcal{E}_{M_k}^F(M_i, M)_0^{co}$: This follows from Corollary 6.1.
- (b) $\mathcal{E}_{M_k}^F(M_i, M)_0^{co} = \cup_{j > i} \mathcal{U}_{k,j}^i$: By Lemma 6.1 (2) for each $f \in \mathcal{E}_{M_k}^F(M_i, M)_0^{co}$ there exists a compact n -submanifold N_1 of M such that $\mathcal{E}_{M_k}^F(M_i, N_1)_0^{co}$ is a neighborhood of f in $\mathcal{E}_{M_k}^F(M_i, M)_0^{co}$. If we choose a $j > i$ such that $N_1 \subset M_j$, then we have $f \in \mathcal{U}_{k,j}^i$.
- (c) $cl \mathcal{U}_{k,j}^i \subset \mathcal{U}_{k,j+1}^i$: Given any $f \in cl \mathcal{U}_{k,j}^i$. Then $f(M_i) \subset M_j \subset U_{j+1}$. First we show that $L \cap U_{j+1}$ is connected for each $L \in \mathcal{C}(M - \text{Int } f(M_i))$. Take $g \in \mathcal{U}_{k,j}^i$ which is sufficiently close to f so that there exists a $k \in G(U_{j+1})$ such that $kg = f$. Since the restriction map $\pi_{k,j}^i : F_{M_k}(U_j)_0 \rightarrow \mathcal{U}_{k,j}^i$ is surjective, there exists an $h \in F_{M_k}(U_j)_0 \subset G(U_{j+1})$ such that $h|_{M_i} = g$. Recall the condition (b) for the exhausting sequence $\{M_i\}_{i \geq 0}$. Then the claim is verified by the homeomorphism of tuples $kh : (M, U_{j+1}, M_i) \approx (M, U_{j+1}, f(M_i))$.

By Lemma 6.1 (1) there exists a neighborhood $\mathcal{V}$ of f in $\mathcal{E}_X^F(N, M)_0^{co}$ and a map $\psi : \mathcal{V} \rightarrow F_{M_k}(U_{j+1})_0$ such that $\psi(g)f = g$ ($g \in \mathcal{V}$) and $\psi(f) = id_M$. Since $\mathcal{V} \cap \mathcal{U}_{k,j}^i \neq \emptyset$, we can choose

a $g' \in \mathcal{V} \cap \mathcal{U}_{k,j}^i$. There exists an $h' \in F_{M_k}(U_j)_0 \subset F_{M_k}(U_{j+1})_0$ such that $h'|_{M_i} = g'$. It follows that $\psi(g')^{-1}h' \in F_{M_k}(U_{j+1})_0$ and $f = \psi(g')^{-1}h'|_{M_i} \in \mathcal{U}_{k,j+1}^i$ as required. $\square$

Remark 6.2. The tuple $(H_X, M, \{M_i\}_{i \geq 1})$ does not satisfy the assumption (A-2) (ii).

Lemma 3.5 and Theorem 3.1 (2)(ii) induce the following conclusions.

Proposition 6.3. (1) If $F_X(U_j)_0 \cap F_{M_i} \simeq *$ for each $j > i \geq 1$, then $(F_X)_0^{c*}$ is HD in $(F_X)_0$.
(2) If F_{M_1} is connected and $F_{M_1}(U_j)_0 \cap F_{M_i} \simeq *$ for each $j > i \geq 2$, then $(F_X)_0^{c*}$ is HD in $(F_X)_0$.

For n -manifolds of finite type, Proposition 4.2 and Proposition 6.3 induce the following conclusions.

Proposition 6.4. Suppose $M = \text{Int } N$ for some compact connected orientable n -manifold N with non-empty boundary, ω is a volume form on M and X is a compact smooth n -submanifold of M . Then the following hold.

- (1) Both $\mathcal{D}_X(M; \omega)_0$ and $(F_X)_0$ are ℓ^2 -manifolds.
- (2) If $\mathcal{D}_{\partial N \times \{0,1\}}(\partial N \times [0,1])_0 \simeq *$, then $(F_X)_0^{c*}$ is HD in $(F_X)_0$.

Proof. (1) The group $(G_X)_0$ is an ANR by Proposition 4.2. Hence, so is $(H_X)_0$ by Proposition 6.1. We can apply Proposition 6.1 (2)(ii) to each $L \in \mathcal{C}(M - \text{Int } X)$ to show that $(F_X)_0$ is a SDR of $(H_X)_0$. Hence $(F_X)_0$ is also an ANR.

(2) We construct an exhausting sequence $\{M_i\}_{i \geq 1}$ of (M, X) as in the proof of Proposition 4.2. Let $U_i = \text{Int } M_i$ ($i \geq 1$). Since this exhausting sequence satisfies the above conditions (a), (b), the tuple $(F_X, M, \{M_i\}_{i \geq 1})$ satisfies the assumption (A). Note that each $L \in \mathcal{C}(M - \text{Int } M_1)$ is a product end.

By Proposition 6.4 (2) it suffices to show the following statements.

- (i) F_{M_1} is connected : By Proposition 6.1 the subgroup H_{M_1} is a SDR of $G_{M_1} = \mathcal{D}_{M_1}^+(M, E_M^\omega)$. We can apply Proposition 6.2 (2)(i) to each $L \in \mathcal{C}(M - \text{Int } M_1)$ to show that F_{M_1} is a SDR of H_{M_1} . Since $G_{M_1} \simeq *$ by the Alexander trick towards ∞ , we have $F_{M_1} \simeq *$.
- (ii) $F_{M_1}(U_j)_0 \cap F_{M_i} \simeq *$ for $j > i \geq 2$: In the proof of Proposition 4.2 (3) we have already shown that $G_{M_1}(U_j)_0 \cap G_{M_i} \simeq *$ for $j > i \geq 2$. By Proposition 6.1 (1) $F_{M_i}(U_j) = H_{M_i}(U_j)$ is a SDR of $G_{M_i}(U_j)$. Then, it follows that $F_{M_1}(U_j)_0 \cap F_{M_i} = F_{M_i}(U_j)_0$ and it is a SDR of $G_{M_1}(U_j)_0 \cap G_{M_i} = G_{M_i}(U_j)_0$. This implies the assertion. $\square$

6.4. Groups of volume-preserving diffeomorphisms of noncompact 2-manifolds.

Suppose M is a noncompact connected orientable smooth 2-manifold without boundary and ω is a volume form on M .

Proof of Theorem 1.3. Since $\mathcal{D}(M; \omega)_0$ is a SDR of $\mathcal{D}(M)_0$ ([31, Corollary 1.1 (2)(ii)], see Section 6.1), the assertions follow from Theorem 1.1 and the observations in Section 6.1. $\square$

Proof of Theorem 1.4. (1) Since $\ker c_0^\omega$ is a SDR of $\mathcal{D}(M; \omega)_0$ ([31, Corollary 1.2 (2)(ii)], see Section 6.1), the assertion follows from Theorem 1.3 and the observations in Section 6.1.

(2) Let $(G, H, F) = (\mathcal{D}(M), \mathcal{D}(M; \omega), \ker c^\omega)$. We have to show that F_0^{c*} is HD in F_0 . We separate the next two cases (cf. Theorem 5.3):

(I) M = a plane or an open annulus. (II) M is not Case (I).

Case (I): Since $M = \text{Int } N$ ($N = \mathbb{D}$ or $\mathbb{A}$), the assertion follows from Proposition 6.4 (2) (cf.)

Case (II): There exists an exhausting sequence $\{M_i\}_{i \geq 1}$ of M which satisfies the conditions (a), (b) in Section 6.3 and such that each M_i is neither a disk, an annulus nor a Möbius band. Let $U_i = \text{int } M_i$ ($i \geq 1$). In the proof of Theorem 5.3, we have shown that

$$G(U_j)_0 \cap G_{M_i} = G_{M_i}(U_j)_0 \simeq * \quad \text{for each } j > i \geq 1.$$

Since $H_{M_i}(U_j)$ is a SDR of $G_{M_i}(U_j)$, this implies that

$$F(U_j)_0 \cap F_{M_i} = H(U_j)_0 \cap H_{M_i} = H_{M_i}(U_j)_0 \simeq * \quad \text{for each } j > i \geq 1.$$

Hence, the conclusion follows from Proposition 6.3. □

REFERENCES

- [1] S. R. Alpern and V.S.Prasad, Typical dynamics of volume-preserving homeomorphisms, Cambridge Tracts in Mathematics, Cambridge University Press (2001).
- [2] T. Banach, K. Mine, K. Sakai and T. Yagasaki, *Homeomorphism and diffeomorphism groups of non-compact manifolds with the Whitney topology*, preprint (arXiv math.GT/0802.0337).
- [3] A. Banyaga, Formes-volume sur les variétés à bord, *Enseignement Math.* (2) 20 (1974) 127 - 131.
- [4] A. Banyaga, The Structure of Classical Diffeomorphism groups, *Mathematics and Its Applications* 400, Kluwer Academic Publishers Group, Dordrecht, 1997.
- [5] R. Berlanga, Groups of measure-preserving homeomorphisms as deformation retracts, *J. London Math. Soc.* (2) 68 (2003) 241 - 254.
- [6] C. Bessaga and A. Pełczyński, Selected topics in infinite-dimensional topology, Polska Akademia Nauk Instytut Mate., Monografie Mate., 58, PWN Polish Scientific Publishers, Warszawa, 1975.
- [7] J. Cerf, *Topologie de certains espaces de plongements*, Bull. Soc. Math. France, 89 (1961) 227–380.
- [8] T. Dobrowolski and H. Toruńczyk, *Separable complete ANR's admitting a group structure are Hilbert manifolds*, Topology Appl., 12 (1981) 229–235.
- [9] C. J. Earle and J. Eells, *A fiber bundle discription of Teichmüller theory*, J. Diff. Geom., 3 (1969) 19–43.
- [10] D. B. A. Epstein, *Curves on 2-manifolds and isotopies*, Acta Math., 155 (1966) 83–107.
- [11] R. Geoghegan, *On spaces of homeomorphisms, embeddings, and functions - I*, Topology, 11 (1972) 159–177.
- [12] R. E. Greene and K. Shiohama, Diffeomorphisms and volume-preserving embeddings of noncompact manifolds, *Trans. Amer. Math. Soc.*, 255 (1979) 403 - 414.
- [13] R. S. Hamilton, *The inverse function theorem of Nash and Moser*, Bull. Amer. Math. Soc. (New Series), 7 n.1 (1982) 65–222.
- [14] M. E. Hamstrom, *Homotopy groups of the space of homeomorphisms on a 2-manifold*, Illinois J. Math., 10 (1966) 563 - 573.
- [15] O. Hanner, *Some theorems on absolute neighborhood retracts*, Ark. Mat., 1 (1951) 389–408.
- [16] M. W. Hirsch, Differential Topology, GTM 33, Springer-Verlag, New York, 1976.
- [17] S. T. Hu, Theory of Retracts, Wayne State Univ. Press, Detroit, 1965.
- [18] J. Kalliongis and D. McCullough, *Fiber-preserving imbeddings and diffeomorphisms*, preprint.
- [19] J. A. Leslie, *On a differential structure for the group of diffeomorphisms*, Topology 6 (1967), 263–271.
- [20] R. Luke and W. K. Mason, *The space of homeomorphisms on a compact two - manifold is an absolute neighborhood retract*, Trans. Amer. Math. Soc., 164 (1972), 275 - 285.
- [21] J. Moser, *On the volume elements on a manifold*, Trans. Amer. Math. Soc., 120 (1965) 286 - 294.
- [22] R. S. Palais, *Local triviality of the restriction map for embeddings*, Comment Math. Helv., 34 (1960) 305–312.
- [23] ———, *Homotopy theory of infinite dimensional manifolds*, Topology, 5 (1966) 1–16.
- [24] G. P. Scott, *The space of homeomorphisms of 2-manifold*, Topology, 9 (1970) 97–109.
- [25] R. T. Seeley, *Extension of C^∞ functions defined in a half space*, Proc. Amer. Math. Soc. 15 (1964), 625–626.
- [26] S. Smale, *Diffeomorphisms of the 2-sphere*, Proc. Amer. Math. Soc., 10 (1959) 621–626.
- [27] H. Toruńczyk, *Characterizing Hilbert space topology*, Fund. Math., 111 (1981) 247–262.
- [28] T. Yagasaki, *Spaces of embeddings of compact polyhedra into 2-manifolds*, Topology Appl., 108 (2000) 107–122.
- [29] ———, *Homotopy types of homeomorphism groups of noncompact 2-manifolds*, Topology Appl., 108 (2000) 123–136.
- [30] ———, *The groups of PL and Lipschitz homeomorphisms of noncompact 2-manifolds*, Bulletin of the Polish Academy of Sciences, Mathematics, 51 (2003) 445–466.

- [31] ———, *Groups of volume-preserving diffeomorphisms of noncompact manifolds and mass flow toward ends*, preprint (arXiv math.GT/0805.3552).

DIVISION OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE AND TECHNOLOGY, KYOTO INSTITUTE OF TECHNOLOGY, MATSUGASAKI, SAKYOKU, KYOTO 606-8585, JAPAN
E-mail address: `yagasaki@kit.ac.jp`